

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2007 (22.02.2007)

PCT

(10) International Publication Number
WO 2007/021991 A2

(51) International Patent Classification:

A61K 31/496 (2006.01) C07D 403/02 (2006.01)
A61K 31/495 (2006.01)

(21) International Application Number:

PCT/US2006/031474

(22) International Filing Date: 11 August 2006 (11.08.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/707,488 11 August 2005 (11.08.2005) US

(71) Applicant (for all designated States except US):
PALATIN TECHNOLOGIES, INC. [US/US]; 4-C
Cedar Brook Drive, Cedar Brook Corporate Center, Cran-
bury, NJ 08512 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SHARMA, Shubh,**
D. [US/US]; 6 Petty Road, Cranbury, NJ 08512 (US). **SHI,**
Yi-Qun [US/US]; 18 Eldrich Street, East Brunswick, NJ
08516 (US). **BURRIS, Kevin** [US/US]; 107 Glenwood
Drive, Washington Crossing, PA 18977 (US). **WU, Zhijun**
[US/US]; 71 Marion Drive, Plainsboro, NJ 08536 (US).
PURMA, Papireddy [IN/US]; 9219 Tamarron Drive,
East Brunswick, NJ 08536 (US). **BONUGA, Yadi, Reddy**
[IN/US]; 9219 Tamarron Drive, Plainsboro, NJ 08536
(US).

(74) Agent: **SLUSHER, Stephen, A.**; Palatin Technologies,
Inc., 4-Cedar Brook Drive, Cranbury, NJ 08512 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT,
LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA,
NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC,
SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ,
UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:


— of inventorship (Rule 4.17(iv))

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: MELANOCORTIN RECEPTOR-SPECIFIC PIPERAZINE AND KETO-PIPERAZINE COMPOUNDS



(57) Abstract: Melanocortin receptor-specific compounds of the general formula (I) and pharmaceu-
tically acceptable salts thereof, where X, W, J, Q, L₁, L₂, L₃, R_{1a}, R_{1b}, R_{2a}, and R_{2b} are as defined in the
specification, and the carbon atom marked with an asterisk can have any stereochemical configuration.
Compounds disclosed herein bind to one or more melanocortin receptors and may be an agonist, a par-
tial agonist, an antagonist, an inverse agonist or an antagonist of an inverse agonist as to one or more
melanocortin receptors, and may be employed for treatment of one or more melanocortin receptor-as-
sociated conditions or disorders, including specifically treatment of obesity and related conditions.

Melanocortin Receptor-Specific Piperazine and Keto-Piperazine Compounds

5

BACKGROUND OF THE INVENTION

Field of the Invention (Technical Field):

The present invention relates to tetra-substituted piperazine compounds that bind to one or more melanocortin receptors and are agonists, antagonists, mixed agonist-
10 antagonists, inverse agonist or antagonists of inverse agonists with respect to one or more melanocortin receptors, and use thereof for the treatment of metabolic, immune, infection-related and other melanocortin receptor-mediated disorders, including treatment of obesity and related energy homeostasis disorders and diseases.

Background Art:

15 A family of melanocortin receptor types and subtypes have been identified, including melanocortin-1 receptors (MC1-R) expressed on normal human melanocytes and melanoma cells, melanocortin-2 receptors (MC2-R) for ACTH (adrenocorticotropin) expressed in cells of the adrenal gland, melanocortin-3 and melanocortin-4 receptors (MC3-R and MC4-R), expressed primarily in cells in the hypothalamus, mid-brain and brainstem, and melanocortin-
20 5 receptors (MC5-R), expressed in a wide distribution of tissues.

In general, compounds specific for MC1-R are believed to be useful for treatment of melanoma. Compounds specific for MC3-R or MC4-R are believed to be useful in regulation of energy homeostasis, including use as agents for attenuating food intake and body weight gain, for use in treatment of anorexia, as a weight gain aid, for treatment of obesity, and
25 treatment of other food intake and metabolism-related purposes. Compounds specific for MC3-R and MC4-R can further be used as agents for treatment of sexual dysfunction, including male erectile dysfunction and female sexual dysfunction. Other melanocortin receptor-specific compounds, such as MCR-1 agonists, can be used as tanning agents to increase melanin production in the skin, acting as chemo-preventive agents against harmful

effects of UV solar radiation. Compounds specific for MCR-1 and MCR-3 may further be useful in regulation of inflammatory processes.

There is a significant need for compounds with high specificity for discrete melanocortin receptors, as well as for compounds that are either agonists or antagonists for specific melanocortin receptors. High affinity compounds for melanocortin receptors can be used to exploit varied physiological responses associated with the melanocortin receptors, either as agonists or antagonists. In addition, melanocortin receptors have an effect on the activity of various cytokines, and high affinity compounds for melanocortin receptors can be used to regulate cytokine activity.

There are piperazine and piperidine compounds known, such as those disclosed in WO 02/070511 (Bristol-Myers Squibb Company), WO 02/059095 (Eli Lilly and Company), and WO 00/74679 (Merck & Co., Inc.), asserted to be specific for melanocortin or related receptors. However, in general such compounds have at most two functional substituted groups, have relatively poor affinity and specificity, and are not suitable for use as a drug compound. There is a significant need for compounds with high specificity for discrete receptors, such as melanocortin and other receptors, as well as compounds that are agonists or antagonists for such receptors. High affinity compounds for such receptors can be used to exploit varied physiological responses associated with the receptors, either as agonists or antagonists. There is thus a need for compounds that are more selective, including higher affinity and specificity, and in particular for compounds that have at least three or four biologically active substituted groups. This invention addresses that need.

WO 02/085925, "Melanocortin Receptor Ligands", to The Proctor & Gamble Company, discloses ketopiperazine structures and methods of synthesis thereof, but does not disclose piperazine structures, piperazine structures with four or more substituted groups, methods to synthesize piperazine structures, methods to synthesize piperazine structures with four or more substituted groups, or methods to synthesize optically pure structures, and further does not disclose structures with a single substituent group that is a single D-Phe or D-Nal residue, or a derivative or homolog thereof, optionally with an amine capping group.


Commonly owned U.S. Patent Application Serial No. 10/837,519, published as Publication No. US 2004/0224957 A1, discloses piperazine compounds specific for one or more melanocortin receptors, but does not disclose piperazine or keto-piperazine compounds with four substituted groups where one substituted group includes a heteroatom unit with at least one cationic center, hydrogen bond donor or hydrogen bond acceptor, and the remaining three substituted groups each include a ring structure.

With respect to certain objects, methods, synthetic schemes, utilities, applications, definitions, protocols and other disclosures, this application is related to U.S. Patent Application Serial No. 10/762,079, entitled "Piperazine Melanocortin-Specific Compounds", filed on January 21, 2004 and International Application No. PCT/US02/25574, International Publication No. WO 03/013571, entitled "Peptidomimetics of Biologically Active Metallopeptides", filed on August 12, 2002; and the specifications of each of the foregoing are incorporated herein by reference as if set forth in full.

There remains a significant need for compounds specific for MC4-R for treatment of conditions relating to regulation of energy homeostasis, including use as agents for attenuating food intake and body weight gain, for treatment of obesity, and treatment of other food intake and metabolism-related purposes.

SUMMARY OF THE INVENTION

In one aspect the invention provides a compound having the formula of structure I:



or an enantiomer, stereoisomer or diastereoisomer thereof, or a pharmaceutically acceptable salt thereof,

wherein

J is a ring structure selected from the group consisting of substituted or unsubstituted aromatic carbocyclic rings, substituted or unsubstituted non-aromatic carbocyclic rings, substituted or unsubstituted aromatic fused carbobicyclic ring groups, two substituted or unsubstituted aromatic carbocyclic rings wherein the rings are joined by a bond, $-\text{CH}_2-$, or $-\text{O}-$, and substituted or unsubstituted aromatic fused heterobicyclic ring groups, wherein in each instance the rings include 5 or 6 ring atoms;

W is a heteroatom unit with at least one cationic center, hydrogen bond donor or hydrogen bond acceptor wherein at least one heteroatom is nitrogen or oxygen;


Q is an aromatic carbocyclic ring selected from the group consisting of phenyl, substituted phenyl, naphthyl and substituted naphthyl;

L_1 is a bond or a linker unit comprising from one to eight backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

L_2 is a bond or $-(\text{CH}_2)_z-$;

L_3 is a bond or a linker unit comprising from one to nine backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

R_{1a} , R_{1b} , R_{2a} and R_{2b} are selected such that:




the remaining of R_{2a} and R_{2b} is hydrogen and R_{1a} and R_{1b} are each hydrogen, R_{1a} and R_{1b}

together form $=\text{O}$, or one of R_{1a} and R_{1b} is a C_1 to C_6 aliphatic linear or branched chain and

the remaining of R_{1a} and R_{2b} is hydrogen,

or R_{2a} and R_{2b} are each hydrogen

and one of R_{1a} and R_{1b} are



and the remaining of R_{1a} and R_{1b} is hydrogen;







5

X is CH_2 , $\text{C}=\text{O}$ or $\text{C}=\text{S}$;


z is an index value from 1 to 6; and

y is an index value from 0 to 5;

wherein the carbon atom marked with an asterisk can have any stereochemical configuration. In the compound of structure I, J may be




6



unsubstituted or substituted with one or more ring substituents. Where substituted, J may be substituted with one or more ring substituents independently selected from the group consisting of hydroxyl, halogen, sulfonamide, alkyl, -O-alkyl, aryl, and -O-aryl.

5 In the compound of structure I, Q may be




where R_{3a}, R_{3b} and R_{3c} are optional ring substituents, and when one or more are present, are the same or different and independently hydroxyl, halogen, alkyl, -O-alkyl, aryl or -O-aryl groups. In one aspect, at least one of R_{3a}, R_{3b} or R_{3c} is -CH₃ or -O-CH₃. In another aspect, at

10 least one of R_{3a}, R_{3b} or R_{3c} is -Cl or -CF₃.

In the compound of structure I, -L₃-Q and -L₁-J may be the same, where Q and J are an aromatic carbocyclic ring selected from the group consisting of phenyl, substituted phenyl, naphthyl and substituted naphthyl.

The group W in the compound of structure I may include an amine, amide, alcohol,
15 carboxylic acid, ether, ester, guanidine or urea, more than one of the foregoing, or a

combination of the foregoing. Thus W may be of the formula



where R_4 is

NH,

5

O,

CH_2 , provided that R_5 comprises N or O,

C_6H_5 , provided that R_5 comprises N or O,

$\text{N}(\text{CH}_2)_z$ where $\text{N}(\text{CH}_2)_z$ together with R_5 forms a ring,

$\text{N}((\text{CH}_2)_y-\text{CH}_3)$,

10

$\text{NH}-\text{C}(=\text{O})$,

$\text{NH}-\text{C}(=\text{O})-\text{NH}$,

$\text{C}(=\text{O})$,

$\text{C}(=\text{O})-\text{NH}$,

$\text{C}(=\text{O})-\text{O}$, or

15

$\text{O}-\text{C}(=\text{O})$;

R_5 is

NH_2 ,

OH,

CH_3 , provided that R_4 comprises N or O,

$\text{NH}(\text{CH}_2)_z$ where $\text{NH}(\text{CH}_2)_z$ together with R_4 forms a ring,

$\text{NH}(\text{CH}_2)_y-\text{CH}_3$,

$\text{N}(-(\text{CH}_2)_y-\text{CH}_3)_2$,

















$\text{NH}(\text{CH}_2)_z-\text{NH}_2$,

$\text{NH}(\text{CH}_2)_z-\text{NH}(\text{CH}_2)_y-\text{CH}_3$,

$\text{NH}(\text{CH}_2)_z-\text{N}((\text{CH}_2)_y-\text{CH}_3)_2$,


$\text{N}(-(\text{CH}_2)_y-\text{CH}_3)-\text{C}(=\text{NH})-\text{NH}_2$,

$\text{N}(-(\text{CH}_2)_y-\text{CH}_3)-\text{C}(=\text{N}((\text{CH}_2)_y-\text{CH}_3))-\text{NH}_2$,



where one or more of positions 1 to 5 are a heteroatom


selected from N for position 1 and S, O or NH for positions 2 to 5,



where none, one or two of positions 1 to 5 are a

heteroatom selected from N for position 1 and for the position to which R_6 is

bound if such position does not comprise C, and otherwise S, O or NH,




where at least one bond between adjacent ring atoms is a


double bond, and one or more of positions 1 to 5 are a heteroatom selected

from N for position 1 and any double bond position and otherwise S, O or


NH for positions 2 to 5, provided that not more than one position is S or O,




where at least one bond between adjacent ring atoms is a double bond, and one or more of positions 1 to 5 are optionally a heteroatom selected from N for position 1, the position to which R_6 is bound if such position does not comprise C, and any double bond position and otherwise S, O or NH for positions 2 to 5, provided that not more than one position is S or O,




where at least one bond between adjacent ring atoms is a double bond, the oxo is bound to a ring carbon, and one or more of the remaining of positions 1 to 5 are optionally a heteroatom selected from N for position 1 and any double bond position and otherwise S, O or NH for positions 2 to 5, provided that not more than one position is S or O,



where at least one bond between adjacent ring atoms is a double bond, the oxo is bound to a ring carbon, and one or more of positions 1 to 5 are optionally a heteroatom selected from N for position 1, the position to which R_6 is bound if such position does not comprise C, and any double bond position and otherwise S, O or NH for positions 2 to 5, provided that not more than one position is S or O,




where one or more of positions 1 to 6 are a heteroatom selected from N for position 1 and S, O or NH for positions 2 to 6,




where none, one or two of positions 1 to 6 are a


heteroatom selected from N for position 1 and the position to which R_6 is bound if such position does not comprise C, and otherwise S, O or NH,




where at least one bond between adjacent ring atoms is a double bond, and one or more of positions 1 to 6 are a heteroatom selected from N for position 1 and any double bond position and otherwise S, O or NH for positions 2 to 6, provided that not more than two positions are S or O,



where at least one bond between adjacent ring atoms is a double bond, and one or more of positions 1 to 6 are optionally a heteroatom selected from N for position 1, the position to which R_6 is bound if such position does not comprise C, and any double bond position and otherwise S, O or NH for positions 2 to 6, provided that not more than two positions are S or O,



where at least one bond between adjacent ring atoms is a double bond, the oxo is bound to a ring carbon, and one or more of the remaining of positions 1 to 6 are optionally a heteroatom selected from N for position 1 and any double bond position and otherwise S, O or NH for positions 2 to 6, provided that not more than two positions are S or O, or



where at least one bond between adjacent ring atoms is a double bond, the oxo is bound to a ring carbon, and one or more of positions 1 to 6 are optionally a heteroatom selected from N for position 1, the position to which R_6 is bound if such position does not comprise C, and

any double bond position and otherwise S, O or NH for positions 2 to 6,
provided that not more than two positions are S or O;

R_8 is hydroxyl, $(CH_2)_y-CH_3$, $(CH_2)_y-NH_2$, $NH-(CH_2)_y-CH_3$ or $N(-(CH_2)_y-CH_3)_2$;


t is an index value from 0 to 5;

z is an index value from 1 to 6; and

5 y is in each instance independently an index value from 0 to 5;

provided that, any NH or NH_2 in the foregoing may be substituted by N-Prg or NH-Prg, respectively, where each Prg is independently an amine protecting group. In the foregoing depictions of ring structures containing a circle within the ring, it is to be understood that the ring structure may include only one double bond, or may include more than one
10 double bond, and in particular, the use of the circle does not imply that all possible double bonds are present. Each Prg may independently be acetyl, adamantyloxy, benzoyl, benzyl, benzyloxycarbonyl, t-butoxycarbonyl, mesitylene-2-sulfonyl, 4-methoxy-2,3,6-trimethyl-benzenesulfonyl, 2,2,4,6,7-pentamethyldihydrobenzofurane-5-sulfonyl, 2,2,5,7,8-pentamethylchromane-6-sulfonyl, 9-fluorenylmethyloxycarbonyl, or tosyl.

In another aspect, the compound of structure I has the formula:



wherein:

R₇ is H or =O;

5 R₈ is hydrogen or N(R_{9a}R_{9b});


R_{9a} and R_{9b} are each independently each independently hydrogen, acetyl, methyl, ethyl, propyl, isopropyl, butyl, pentyl, hexyl, isobutyl, benzyl, benzoyl, hexanoyl, propionyl, butanoyl, pentanoyl, heptanoyl, cyclopropyl, cyclopropylmethyl, cyclobutyl, cyclobutylmethyl, cyclohexyl, cyclohexylmethyl, or polyethylene glycol;

10 v is in each instance independently an index value from 0 to 2; and


y is in each instance independently an index value from 0 to 5;

wherein if R₈ is not hydrogen, the adjacent carbon atom marked with an asterisk can have any stereochemical configuration. Where R_{9a} or R_{9b} are polyethylene glycol, the polyethylene glycol has a formula molecular weight of between 100 and 50,000.

In one aspect of the compound of structure I, one of R_{2a} and R_{2b} is




or



and the remaining of R_{2a} and R_{2b}

and both R_{1a} and R_{1b} are hydrogen.

The invention further provides a compound having the formula of structure II:



or an enantiomer, stereoisomer or diastereoisomer thereof, or a pharmaceutically acceptable salt thereof,

wherein

each J is independently a ring structure selected from the group consisting of substituted or unsubstituted aromatic carbocyclic rings, substituted or unsubstituted non-aromatic carbocyclic rings, substituted or unsubstituted aromatic fused carbobicyclic ring groups, two substituted or unsubstituted aromatic carbocyclic rings wherein the rings are joined by a bond, $-CH_2-$, or $-O-$, and substituted or unsubstituted aromatic fused heterobicyclic ring groups, wherein in each instance the rings include 5 or 6 ring atoms;


W is a heteroatom unit with at least one cationic center, hydrogen bond donor or hydrogen bond acceptor wherein at least one heteroatom is nitrogen or oxygen;

L_1 is a bond or a linker unit comprising from one to eight backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

L_2 is a bond or $-(CH_2)_z-$;

L_3 is a bond or a linker unit comprising from one to eight backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

R_{1a} , R_{1b} , R_{2a} and R_{2b} are selected such that:




the remaining of R_{2a} and R_{2b} is hydrogen and R_{1a} and R_{1b} are each hydrogen, R_{1a} and R_{1b} together form $=O$, or one of R_{1a} and R_{1b} is a C_1 to C_6 aliphatic linear or branched chain and the remaining of R_{1a} and R_{2b} is hydrogen,

or R_{2a} and R_{2b} are each hydrogen

5

and one of R_{1a} and R_{1b} are



or

,

and the remaining of R_{1a} and R_{1b} is hydrogen;


X is CH_2 , $C=O$ or $C=S$;

z is an index value from 1 to 6; and


y is an index value from 0 to 5;

10


wherein the carbon atom marked with an asterisk can have any stereochemical configuration. In the compounds of structure II, each J may independently be:




,




,




,




,



,



,



unsubstituted or substituted with one or more ring substituents, such substituents independently selected from the group consisting of hydroxyl, halogen, sulfonamide, alkyl, -O-alkyl, aryl, and -O-aryl. In one particular aspect of the compound of structure II, -L₃-J and -L₁-J are the same.

The invention further provides a pharmaceutical composition comprising a compound of structure I or structure II and a pharmaceutically acceptable carrier. Such pharmaceutical composition may be employed for affecting melanocortin receptor function in a human or non-human mammal. Such pharmaceutical composition may further be employed for treating a condition responsive to changes in melanocortin receptor function in a human or non-human mammal, by administering the pharmaceutical composition to the human or non-human mammal in a pharmaceutically effective amount. The condition may be selected from the group consisting of male sexual dysfunction, female sexual dysfunction, an eating disorder, above-optimal body weight, obesity, below-optimal body weight and cachexia.

The present invention further provides compounds that are agonists of a melanocortin receptor, including one or more of MC1-R, MC3-R, MC4-R, or MC5-R. The compounds

alternatively are antagonists of a melanocortin receptor, including one or more of MC1-R, MC3-R, MC4-R, or MC5-R. The compounds alternatively are inverse agonists of a melanocortin receptor, including one or more of MC1-R, MC3-R, MC4-R, or MC5-R. The compounds alternatively are antagonists of an inverse agonist of a melanocortin receptor, including one or more of MC1-R, MC3-R, MC4-R, or MC5-R.

The invention further includes methods for altering a disorder or condition associated with the activity of a melanocortin receptor, comprising administering to a patient a pharmaceutically effective amount a compound of this invention. In one embodiment the disorder or condition is an eating disorder such as cachexia. In another embodiment the disorder or condition is obesity and associated impairment of energy homeostasis. In yet another embodiment the disorder or condition is sexual dysfunction such as erectile dysfunction or female sexual dysfunction.

One object of the present invention is to provide conformationally constrained and optically pure isomers of tetra-substituted piperazine, wherein the pendant group substituents are amino acid moieties, amino acid side chain moieties or derivatives thereof, such that the resulting ring compound biologically mimics a relevant reverse turn peptide structure.

Another object of the present invention is to provide methods for the synthesis of optically pure tetra-substituted piperazine compounds.

Another object of the present invention is to provide piperazine compounds with four pendant groups, such pendant groups consisting of any moiety other than H, O, S, or a halogen.

Another object of the present invention is to provide piperazine core compounds wherein pendant groups are provided, which pendant groups are or include amino acid side chain moieties.

Another object of the present invention is to provide a tetra-substituted piperazine compound wherein such compound is specific for one or more melanocortin receptors.

Another object of the present invention is to provide a method for synthesis of tetra-substituted piperazine compounds of the invention.

Other objects, advantages and novel features, and the further scope of applicability of


the present invention will be set forth in part in the detailed description to follow, and in part will become apparent to those skilled in the art upon examination of the following, or may be learned by practice of the invention.

5

DETAILED DESCRIPTION OF THE INVENTION

In this invention it is disclosed that piperazine and keto-piperazine rings may be employed with four descriptors, wherein each descriptor is a separate pendant group covalently bound to a given ring atom, and where at least three descriptors include a ring structure. The rings in the ring structure in each of the descriptors or pendant groups may be heterocyclic or carbocyclic, and at least two of such rings are aromatic. By employing four descriptors, the inventors have further found that the chirality of the ring, and stereo structure generally, is fixed in a desired structure, thereby more closely mimicking the desired pharmacophores, and with the descriptors positioned in the most relevant chemical space.


The application includes keto-piperazine as well as piperazine compounds. Thus the following compounds, by way of example, are included within the invention:



20

where L_1 , L_2 , L_3 , J , W , Q and y are as defined in the Summary of the Invention.

The invention further includes compounds which are penta-substituted piperazine or keto-piperazine compounds, wherein the fifth substituent is a C_1 to C_6 aliphatic linear or branched chain. The following are representative examples of this class, it being understood
 5 that the methyl group shown may be any C_1 to C_6 aliphatic linear or branched chain:



where here also L_1 , L_2 , L_3 , J , W , Q and y are as defined in the Summary of the Invention.

This invention thus discloses the use of tetra- and penta-substituted piperazine and
 10 keto-piperazine templates for drug design where four substituents are provided, at least three of which include a ring structure. The invention further also relates to enantiomerically pure compounds, preferably made by the synthetic schemes disclosed herein or variants thereof. A classical piperazine ring is a conformationally dynamic six-membered ring structure. It can exist in a variety of conformational states, commonly referred to as chair, boat, twisted chair
 15 or twisted boat conformations. Because of this dynamism in structural states, the location of descriptors on the ring plays an important role in stabilizing the ring in a single conformational state; if the appropriate conformational state is selected, this is conducive to making a molecule more selective for its receptor. For example, a 1,3 axial placement of two bulky descriptors generally causes unfavorable steric interactions between these two groups, and
 20 thus make a chair conformation energetically less stable. Consequently, the chair conformation is less preferred, resulting in a twisted chair or boat conformation. The twisted chair or boat conformation results in a specific stereochemical alignment of the descriptors, which is specifically relevant to interaction with the desired receptor. Thus, a conformation

resulting from 1,3 axial placement of two descriptors may result in a structure more selective for a given receptor sub-type.

In yet another embodiment, the invention describes piperazine and keto-piperazine compounds specific for G-protein coupled receptor systems, such systems including, but not
5 limited to, melanotropin or melanocortin receptors (MC1-R, MC3-R, MC4-R and MC5-R).

In yet another embodiment, the invention provides novel schemes and methods of synthesis of piperazine and keto-piperazine compounds.

Definitions. Before proceeding further with the description of the invention, certain terms are defined as set forth herein.

10 The "amino acid" and "amino acids" used in this invention, and the terms as used in the specification and claims, include the known naturally occurring protein amino acids, which are referred to by both their common three letter abbreviation and single letter abbreviations. See generally Synthetic Peptides: A User's Guide, GA Grant, editor, W.H. Freeman & Co., New York, 1992, the teachings of which are incorporated herein by reference, including the
15 text and table set forth at pages 11 through 24. As set forth above, the term "amino acid" also includes stereoisomers and modifications of naturally occurring protein amino acids, non-protein amino acids, post-translationally modified amino acids, enzymatically synthesized amino acids, derivatized amino acids, constructs or structures designed to mimic amino acids, and the like. Modified and unusual amino acids are described generally in Synthetic
20 Peptides: A User's Guide, cited above; Hruby VJ, Al-obeidi F and Kazmierski W: *Biochem J* 268:249-262, 1990; and Toniolo C: *Int J Peptide Protein Res* 35:287-300, 1990; the teachings of all of which are incorporated herein by reference.

The term "amino acid side chain moiety" used in this invention includes any side chain of any amino acid, as the term "amino acid" is defined herein, including any derivative of
25 an amino acid side chain moiety, as the term "derivative" is defined herein. This thus includes the side chain moiety present in naturally occurring amino acids. It further includes side chain moieties in modified naturally occurring amino acids, such as glycosylated amino acids. It further includes side chain moieties in stereoisomers and modifications of naturally occurring protein amino acids, non-protein amino acids, post-translationally modified amino acids,

enzymatically synthesized amino acids, derivatized amino acids, constructs or structures designed to mimic amino acids, and the like. For example, the side chain moiety of any amino acid disclosed herein is included within the definition of an amino acid side chain moiety.

5 The "derivative" of an amino acid side chain moiety includes any modification to or variation in any amino acid side chain moieties, including a modification of naturally occurring amino acid side chain moieties. By way of example, derivatives of amino acid side chain moieties include straight chain or branched, cyclic or noncyclic, substituted or unsubstituted, and saturated or unsaturated alkyl, aryl or aralkyl moieties.

10 Conventional amino acid residue abbreviations have their conventional meaning as given in Chapter 2400 of the Manual of Patent Examining Procedure, 8th Ed. Thus, "Nle" is norleucine; "Asp" is aspartic acid; "His" is histidine; "D-Phe" is D-phenylalanine; "Arg" is arginine; "Trp" is tryptophan; "Lys" is lysine; "Gly" is glycine; "Pro" is proline; "Tyr" is tyrosine, "Ser" is serine and so on.

15 In the specification and the claims, the term "homolog" includes, without limitation, (a) a D-amino acid residue or side chain substituted for an L-amino acid residue side chain, (b) a post-translationally modified residue or side chain substituted for the residue or side chain, (c) a non-peptide or other modified amino acid residue or side chain based on another such residue or side chain, such as phenylglycine, homophenylalanine, ring-substituted
20 halogenated, and alkylated or arylated phenylalanines for a phenylalanine residue, diamino proionic acid, diamino butyric acid, ornithine, lysine and homoarginine for an arginine residue, and the like, and (d) any amino acid residue or side chain, coded or otherwise, or a construct or structure that mimics an amino acid residue or side chain, and which has at least a
25 similarly charged side chain (neutral, positive or negative), preferably a similar hydrophobicity or hydrophilicity, and preferably a similar side chain in terms of being a saturated aliphatic side chain, a functionalized aliphatic side chain, an aromatic side chain or a heteroaromatic side chain.

 The term "alkene" includes unsaturated hydrocarbons that contain one or more double carbon-carbon bonds. Examples of such alkene groups include ethylene, propene,

and the like.

The term "alkenyl" includes a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms containing at least one double bond; examples thereof include ethenyl, 2-propenyl, and the like.

The "alkyl" groups specified herein include those alkyl radicals in either a straight or branched configuration. Examples of such alkyl radicals include methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tertiary butyl, pentyl, isopentyl, hexyl, isohexyl, and the like.

The term "alkynal" includes a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms containing at least one triple bond; examples thereof include ethynyl, propynal, butynyl, and the like.

The term "aryl" includes a monocyclic or bicyclic aromatic hydrocarbon radical of 6 to 12 ring atoms, and optionally substituted independently with one or more substituents selected from alkyl, haloalkyl, cycloalkyl, alkoxy, alkylthio, halo, nitro, acyl, cyano, amino, monosubstituted amino, disubstituted amino, hydroxy, carboxy, or alkoxy-carbonyl. Examples of an aryl group include phenyl, biphenyl, naphthyl, 1-naphthyl, and 2-naphthyl, derivatives thereof, and the like.

The term "aralkyl" includes a radical $-R^aR^b$ where R^a is an alkylene (a bivalent alkyl) group and R^b is an aryl group as defined above. Examples of aralkyl groups include benzyl, phenylethyl, 3-(3-chlorophenyl)-2-methylpentyl, and the like.

The term "aliphatic" includes compounds with hydrocarbon chains, such as for example alkanes, alkenes, alkynes, and derivatives thereof.

The term "acyl" includes a group $RCO-$, where R is an organic group. An example is the acetyl group CH_3CO- .

A group or aliphatic moiety is "acylated" when an alkyl or substituted alkyl group as defined above is bonded through one or more carbonyl $[-(C=O)-]$ groups.

An "omega amino derivative" includes an aliphatic moiety with a terminal amino group. Examples of omega amino derivatives include aminoheptanoyl and the amino acid

side chain moieties of ornithine and lysine.

The term "heteroaryl" includes mono- and bicyclic aromatic rings containing from 1 to 4 heteroatoms selected from nitrogen, oxygen and sulfur. 5- or 6-membered heteroaryl are monocyclic heteroaromatic rings; examples thereof include thiazole, oxazole, thiophene, furan, pyrrole, imidazole, isoxazole, pyrazole, triazole, thiadiazole, tetrazole, oxadiazole, pyridine, pyridazine, pyrimidine, pyrazine, and the like. Bicyclic heteroaromatic rings include, but are not limited to, benzothiadiazole, indole, benzothiophene, benzofuran, benzimidazole, benzisoxazole, benzothiazole, quinoline, benzotriazole, benzoxazole, isoquinoline, purine, furopyridine and thienopyridine.

An "amide" includes compounds that have a trivalent nitrogen attached to a carbonyl group ($-\text{CO.NH}_2$), such as methylamide, ethylamide, propylamide, and the like.

An "imide" includes compounds containing an imido group ($-\text{CO.NH.CO}-$).

An "amine" includes compounds that contain an amino group ($-\text{NH}_2$).

A "nitrile" includes compounds that are carboxylic acid derivatives and contain a ($-\text{CN}$) group bound to an organic group.

An amino acid side chain moiety is "hydrogen bonding" when the side chain includes hydrogen bond donors and/or hydrogen bond acceptors.

An "amine capping group" includes any terminal group attached through a terminal amine, including but not limited to any omega amino derivative, acyl group or terminal aryl or aralkyl, including groups such as a C_1 to C_6 linear or branched chain such as methyl, dimethyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, or hexyl, groups such as allyl, cyclopropane methyl, hexanoyl, heptanoyl, acetyl, propionoyl, butanoyl, phenylacetyl, cyclohexylacetyl, naphthylacetyl, cinnamoyl, phenyl, benzyl, benzoyl, 12-Ado, 7'-amino heptanoyl, 6-Ahx, Amc or 8-Aoc, or a molecule such as polyethylene glycol with a formula molecular weight of between 100 and 50,000.

The term "composition", as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one

or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing a compound of the present invention and one or more pharmaceutically acceptable carriers and/or other excipients, and optionally one or more other pharmaceutically active ingredients and agents.

A variety of chemicals and compounds are employed in this invention, and the following abbreviations have the meanings given:

	Boc	tertiary butyloxycarbonyl
	Cbz	benzyloxycarbonyl
10	DCM	dichloromethane
	DIAD	diisopropyl azodicarboxylate
	DMF	N,N-dimethylformamide
	DMSO	dimethyl sulfoxide
	EDC	N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride
15	EtOAc	ethyl acetate
	Fmoc	9-fluorenylmethoxycarbonyl
	HEPES	4-(2-hydroxyethyl)1-piperazineethanesulfonic acid
	HOAt	1-hydroxy-7-azabenzotriazole
	IBCF	isobutyl chloroformate
20	LAH	lithium aluminum hydride
	NMM	N-methylmorpholine
	Pd/C	palladium on carbon
	TBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate
25	TEA	triethylamine
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
	TPP	triphenylphosphine

A "tetra-substituted piperazine", as used herein, is a piperazine compound or

derivative thereof wherein a group other than solely H, and preferably including an amino acid residue or an amino acid side chain moiety, is attached to each ring N member, and further wherein groups other than solely H, O, S or a halogen, preferably including an amino acid side chain moiety, are attached to two ring C members.

5 "Sexual dysfunction" means any condition that inhibits or impairs normal sexual function, including coitus. The term is not limited to physiological conditions, and includes psychogenic conditions or perceived impairment without a formal diagnosis of pathology or disorder. Sexual dysfunction includes erectile dysfunction in a male mammal and female sexual dysfunction in a female mammal.

10 "Erectile dysfunction" is a disorder involving the failure of a male mammal to achieve functional erection, ejaculation, or both. Erectile dysfunction is accordingly synonymous with impotence, and may include the inability to attain or sustain an erection of sufficient rigidity for coitus. Symptoms of erectile dysfunction include an inability to achieve or maintain an erection, ejaculatory failure, premature ejaculation, or inability to achieve an orgasm, which
15 symptoms may occur separately or in any combination. An increase in erectile dysfunction is often associated with age or may be caused by a physical disease or as a side-effect of drug treatment.

 "Female sexual dysfunction" is a disorder including sexual arousal disorder. The term "sexual arousal disorder" includes a persistent or recurrent failure to attain or maintain the
20 lubrication-swelling response of sexual excitement until completion of sexual activity. Sexual dysfunction in females may also include inhibited orgasm and dyspareunia, which is painful or difficult coitus. Female sexual dysfunction includes, but is not limited to, a number of categories of diseases, conditions and disorders including hypoactive sexual desire disorder, sexual anhedonia, sexual arousal disorder, dyspareunia and vaginismus. Hypoactive sexual
25 desire disorder includes a disorder in which sexual fantasies and desire for sexual activity are persistently or recurrently diminished or absent, causing marked distress or interpersonal difficulties. Hypoactive sexual desire disorder can be caused by boredom or unhappiness in a long-standing relationship, depression, stress, dependence on alcohol or psychoactive drugs, side effects from prescription drugs, or hormonal deficiencies. Sexual anhedonia

includes decreased or absent pleasure in sexual activity. Sexual anhedonia can be caused by depression, drugs, or interpersonal factors. Sexual arousal disorder can be caused by reduced estrogen, illness, or treatment with diuretics, antihistamines, antidepressants or antihypertensive agents. Dyspareunia and vaginismus are sexual pain disorders

5 characterized by pain resulting from penetration and may be caused, for example, by medications which reduce lubrication, endometriosis, pelvic inflammatory disease, inflammatory bowel disease or urinary tract problems.

By a melanocortin receptor "agonist" is meant an endogenous or drug substance or compound, including a compound of this invention, which can interact with a melanocortin

10 receptor and initiate a pharmacological response characteristic of the melanocortin receptor. By a melanocortin receptor "antagonist" is meant a drug or a compound, including a compound of this invention, which opposes the melanocortin receptor-associated responses normally induced by a melanocortin receptor agonist agent.

By "binding affinity" is meant the ability of a compound or drug to bind to its biological

15 target.

The chemical naming protocol and structure diagrams used herein employ and rely on the chemical naming features as utilized by the ChemDraw program (available from Cambridgesoft Corp.) or ISIS Draw (MDL Information Systems, Inc.). In particular, the compound names were derived from the structures using the Autonom program as utilized by

20 ChemDraw Ultra or ISIS Draw.

Clinical Applications. The compounds disclosed herein can be used for both medical applications and animal husbandry or veterinary applications. Typically, the product is used in humans, but may also be used in other mammals. The term "patient" is intended to denote a mammalian individual, and is so used throughout the specification and in the claims.

25 The primary applications of this invention involve human patients, but this invention may be applied to laboratory, farm, zoo, wildlife, pet, sport or other animals.

Melanocortin receptor-specific compounds of this invention that are MC1-R specific can be used as chemoprevention agents against sun-induced, such as by UV radiation, neoplastic activity in human skin. MC1-R agonist compounds of this invention may be

employed to stimulate epidermal melanocytes to produce melanin as well as to convert pheomelanin to eumelanin. Eumelanin, which is dark brown or black pigmentation, is considered more photo-protective than pheomelanin, which is yellow or red pigmentation.

The process of melanogenesis is believed to involve stimulation of MC1-R in epidermal

5 melanocytes, thereby mediating the stimulation of tyrosinase enzymes within these pigment cells, inducing the conversion of tyrosine to dopa and then through dopaquinone to

eumelanin. Sun tanning due to direct sun exposure is proposed to result from the same pathway by local production of melanotropic peptide from a POMC gene in the epidermis.

Thus, stimulation of eumelanin production and conversion of pheomelanin to eumelanin may

10 be a desirable chemoprevention modality in blocking sun- or UV-induced neoplastic activity in skin. A potent, high-affinity and highly selective MC1-R agonist compound of this invention can accordingly be used as a therapeutic chemoprevention agent for combating harmful sun or UV exposure that induces neoplastic activity in skin melanocytes.

In another embodiment, compounds of this invention, including but not limited to

15 compounds that are MC4-R agonists, partial agonists or functionally inactive may be used as a therapeutic agent to modify energy metabolism and feeding behavior, including treatment of pathologic obesity and related conditions. In addition to use in treatment of patients clinically diagnosed as obese, compounds of this invention may be employed with persons who are above optimal body weight, as an aid in weight loss. Compounds of this invention, including
20 but not limited to MC4-R antagonists, may be used as a therapeutic agent in eating disorders, such as treatment of anorexia and cachexia, which is malnutrition and wasting due to illness.

In addition to use in treatment of patients diagnosed with anorexia or cachexia, compounds of this invention may be employed with persons who have below optimal body weight, and in particular with patients desiring to gain additional muscle mass.

25 In yet another embodiment, compounds of this invention can be used as therapeutic agents for treatment of sexual dysfunction, including treatment of both male erectile dysfunction and female sexual dysfunction.

In yet another embodiment, compounds of this invention may be used as therapeutic agents for treatment of inflammation, including specifically MC1-R, MC3-R and MC5-R

agonists.

In yet another embodiment of the invention, compounds of this invention that are MC5-R specific can be used as agents to decrease sebum production, and thus may be efficacious in the treatment of acne and related diseases. The compounds for this application may be
5 conveniently formulated for local administration, as through a gel, lotion, cream or other topical formulation.

In yet another embodiment, compounds of this invention may be employed in the treatment of drug or alcohol dependence, depression, anxiety and related conditions and indications.

10 **Formulations and Administration.** The compounds may be formulated by any means, such as those known in the art, including but not limited to tablets, capsules, caplets, suspensions, powders, lyophilized forms and aerosols/aerosolizable formulations and may be mixed and formulated with buffers, binders, stabilizers, anti-oxidants and other agents known in the art. The compounds may be administered by any systemic or partially systemic means
15 such as those known in the art, including but not limited to intravenous injection, subcutaneous injection, administration through mucous membranes, oral administration, dermal administration, skin patches, aerosols and the like.

The invention further provides a pharmaceutical composition that includes a compound of this invention and a pharmaceutically acceptable carrier. The compound of this
20 invention may thus be formulated or compounded into pharmaceutical compositions that include at least one compound of this invention together with one or more pharmaceutically acceptable carriers, including excipients, such as diluents, carriers and the like, and additives, such as stabilizing agents, preservatives, solubilizing agents, buffers and the like, as may be desired. Formulation excipients may include polyvinylpyrrolidone, gelatin, hydroxy cellulose,
25 acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate. For injection or other liquid administration formulations, water containing at least one or more buffering constituents is suitable, and stabilizing agents, preservatives and solubilizing agents may also be employed. For solid administration formulations, any of a variety of thickening, filler, bulking and carrier additives may be employed, such as starches, sugars, fatty acids and the like.

For topical administration formulations, any of a variety of creams, ointments, gels, lotions and the like may be employed. For most pharmaceutical formulations, non-active ingredients will constitute the greater part, by weight or volume, of the preparation. For pharmaceutical formulations, it is also contemplated that any of a variety of measured-release, slow-release
5 or time-release formulations and additives may be employed, such that the dosage may be formulated so as to effect delivery of a compound of this invention over a period of time.

The compounds of this invention may be in the form of any pharmaceutically acceptable salt. Acid addition salts of the compounds of this invention are prepared in a suitable solvent from the compound and an excess of an acid, such as hydrochloric,
10 hydrobromic, sulfuric, phosphoric, acetic, trifluoroacetic, maleic, succinic or methanesulfonic. The acetate salt form is especially useful. Where the compounds of this invention include an acidic moiety, suitable pharmaceutically acceptable salts may include alkali metal salts, such as sodium or potassium salts, or alkaline earth metal salts, such as calcium or magnesium salts.

15 The compounds and pharmaceutical compositions of this invention may be administered by injection, which injection may be intravenous, subcutaneous, intramuscular, intraperitoneal or by any other means known in the art. In general, any route of administration by which the compounds of this invention are introduced across an epidermal layer of cells may be employed. Administration means may include administration through mucous
20 membranes, buccal administration, oral administration, dermal administration, inhalation administration, nasal administration and the like. The dosage for treatment is administration, by any of the foregoing means or any other means, of an amount sufficient to bring about the desired therapeutic effect.

One advantageous route of administration is nasal administration, such as by means
25 of a liquid spray, gel or powder. In one route of administration, an aqueous solution is employed, preferably administered by means of a metered delivery device. By "nasal administration" is meant any form of intranasal administration of any of the compounds and pharmaceutical compositions of this invention. Thus in one embodiment, compounds and pharmaceutical compositions of this invention include an aqueous solution, such as a solution

including saline, citrate or other common excipients or preservatives, formulated for intranasal administration. In another embodiment, compounds and pharmaceutical compositions of this invention include a dry or powder formulation, formulated for intranasal administration. A preparation for nasal administration can take a variety of forms, such as for administration in
5 nasal drops, nasal spray, gel, ointment, cream, powder or suspension. A variety of dispensers and delivery vehicles are known in the art, including single-dose ampoules, metered dose devices, atomizers, nebulizers, pumps, nasal pads, nasal sponges, nasal capsules, and the like.

The pharmaceutical composition may be in a solid, semi-solid, or liquid form. For a
10 solid form, the compound and other components may be mixed together by blending, tumble mixing, freeze-drying, solvent evaporation, co-grinding, spray-drying, and/or other techniques known in the art. A semi-solid pharmaceutical composition suitable for intranasal administration may take the form of an aqueous or oil-based gel or ointment. For example,
the compound and other components can be mixed with microspheres of starch, gelatin,
15 collagen, dextran, polylactide, polyglycolide or other similar materials that form hydrophilic gels. In one embodiment, the microspheres can be internally loaded or coated with compound, which upon administration forms a gel that adheres to the nasal mucosa. In another embodiment, the formulation is liquid, it being understood that this includes, for example, an aqueous solution, an aqueous suspension, an oil solution, an oil suspension, or
20 an emulsion, depending on the physicochemical properties of the compound and other components.

For liquid formulations, excipients necessary or desirable for formulation, stability, and/or bioavailability may be included in the pharmaceutical composition. Exemplary excipients include sugars (such as glucose, sorbitol, mannitol, or sucrose), uptake enhancers
25 (such as chitosan), thickening agents and stability enhancers (such as celluloses, polyvinyl pyrrolidone, starch, and the like), buffers, preservatives, and/or acids and bases to adjust the pH. In one embodiment, an absorption promoting component is included in the pharmaceutical composition. Exemplary absorption promoting components include surfactant acids, such as cholic acid, glycocholic acid, taurocholic acid, and other cholic acid derivatives,

chitosan and cyclodextrins.

The pharmaceutical composition may further include optional components such as humectants, preservatives and the like. A humectant or moisturizing agent can be employed to decrease water loss from the pharmaceutical composition and optionally moisturize nasal mucosa. Exemplary humectants include hygroscopic materials such as glycerine, propylene glycol, polyethylene glycol, polysaccharides and the like. Preservatives may be employed, to prevent or limit bacteria and other microbial growth. One such preservative that may be employed is benzalkonium chloride, such as 0.05% benzalkonium chloride. Other preservatives include, for example, benzyl alcohol, methylparaben, propylparaben, butylparaben, chlorobutanol, phenethyl alcohol, phenyl mercuric acetate and the like.

The pharmaceutical composition may also include rheology modifying agents, such as for varying the viscosity of the pharmaceutical composition. Exemplary rheology modify agents include polyers and similar materials, such as sodium carboxymethyl cellulose, algin, carageenans, carbomers, galactomannans, hydroxypropyl methylcellulose, hydroxypropyl cellulose, polyethylene glycols, polyvinyl alcohol, polyvinylpyrrolidone, sodium carboxymethyl chitin, sodium carboxymethyl dextran, sodium carboxymethyl starch, xanthan gum and combinations of the foregoing. Such agents may also act as bioadhesives, to extend the residence time of a compound of the invention within the nasal mucosa.

Depending on the formulation and route of administration, an aqueous solution of compounds or pharmaceutical compositions of this invention may be appropriately buffered by means of saline, acetate, phosphate, citrate, acetate or other buffering agents, which are at any physiologically acceptable pH, generally from about pH 4 to about pH 8. A combination of buffering agents may also be employed, such as phosphate buffered saline, a saline and acetate buffer, and the like. In the case of saline, a 0.9% saline solution may be employed. In the case of acetate, phosphate, citrate, acetate and the like, a 50 mM solution may be employed.

In another route of administration, compounds and pharmaceutical compositions of this invention are administered directly into the lung. Intrapulmonary administration may be performed by means of a metered dose inhaler, a device allowing self-administration of a

metered bolus of a compound and pharmaceutical composition of this invention when actuated by a patient during inspiration. Both dry powder inhalation and nebulized aerosols may be employed. Thus, it is possible and contemplated that compounds and pharmaceutical compositions of this invention may be in a dried and particulate form. In one

5 embodiment, the particles are between about 0.5 and 6.0 μm , such that the particles have sufficient mass to settle on the lung surface and not be exhaled, but are small enough that they are not deposited on surfaces of the air passages prior to reaching the lung. Any of a variety of different techniques may be used to make dry powder microparticles, including but not limited to micro-milling, spray drying and a quick freeze aerosol followed by lyophilization.

10 With micro-particles, the constructs may be deposited to the deep lung, thereby providing quick and efficient absorption into the bloodstream. Further, with such an approach, penetration enhancers are not required, as is sometimes necessary with transdermal, nasal or oral mucosal delivery routes. Any of a variety of inhalers may be employed, including propellant-based aerosols, nebulizers, single dose dry powder inhalers and multidose dry

15 powder inhalers. Common devices in current use include metered dose inhalers, which are used to deliver medications for the treatment of asthma, chronic obstructive pulmonary disease and the like. Preferred devices include dry powder inhalers, designed to form a cloud or aerosol of fine powder with a particle size that is always less than about 6.0 μm .

Microparticle size, including mean size distribution, may be controlled by means of the

20 method of making. For micro-milling, the size of the milling head, speed of the rotor, time of processing and the like control the microparticle size. For spray drying, the nozzle size, flow rate, dryer heat and the like control the microparticle size. For making by means of quick freeze aerosol followed by lyophilization, the nozzle size, flow rate, concentration of aerosoled solution and the like control the microparticle size. These parameters and others may be

25 employed to control the microparticle size.

The compounds and pharmaceutical compositions of this invention may be formulated for and administered by means of an injection, such as a deep intramuscular injection, such as in the gluteal or deltoid muscle, of a time release injectable formulation. In one embodiment, a compound or pharmaceutical composition of this invention is formulated with a

PEG, such as poly(ethylene glycol) 3350, and optionally one or more additional excipients and preservatives, including but not limited to excipients such as salts, polysorbate 80, sodium hydroxide or hydrochloric acid to adjust pH, and the like. In another embodiment a compound or pharmaceutical composition of this invention is formulated with a poly(ortho ester), which may be an auto-catalyzed poly(ortho ester) with any of a variable percentage of lactic acid in the polymeric backbone, and optionally one or more additional excipients. In one embodiment, poly (D,L-lactide-co-glycolide) polymer (PLGA polymer) is employed, preferably a PLGA polymer with a hydrophilic end group, such as PLGA RG502H from Boehringer Ingelheim, Inc. (Ingelheim, Germany). Such formulations may be made, for example, by combining a compound of this invention in a suitable solvent, such as methanol, with a solution of PLGA in methylene chloride, and adding thereto a continuous phase solution of polyvinyl alcohol under suitable mixing conditions in a reactor. In general, any of a number of injectable and biodegradable polymers, which are preferably also adhesive polymers, may be employed in a time release injectable formulation. The teachings of U.S. Patents 4,938,763, 6,432,438, and 6,673,767, and the biodegradable polymers and methods of formulation disclosed therein, are incorporated herein by reference. The formulation may be such that an injection is required on a weekly, monthly or other periodic basis, depending on the concentration and amount of construct, the biodegradation rate of the polymer, and other factors known to those of skill in the art.

Pharmaceutically Effective Amount. In general, the actual quantity of compound of this invention administered to a patient will vary between fairly wide ranges depending upon the mode of administration, the formulation used, and the response desired. The dosage for treatment is administration, by any of the foregoing means or any other means known in the art, of an amount sufficient to bring about the desired effect. This may readily be determined by one of ordinary skill in the art through means such as pharmacokinetic studies, plasma half-life studies, dose escalation studies, and the like. Thus a pharmaceutically effective amount includes an amount of a compound or pharmaceutical composition of this invention that is sufficient to induce the desired effect.

In general, the compounds of this invention are highly active, with dose responses as

low as 0.01 µg/kg, generally with optimal or peak dose responses between about 0.01 µg/kg and 25 µg/kg, depending on the specific compound and the route of administration. For example, the compound can be administered at 0.01, 0.05, 0.1, 0.5, 1, 5, 10, 50, 100, or 500 µg/kg body weight, depending on specific compound selected, the desired response, the route of administration, the formulation and other factors known to those of skill in the art. Conventional dose response studies and other pharmacological means may be employed to determine the optimal dose for a desired effect with a given compound, given formulation and given route of administration.

Combination Therapy and Weight Regulation. It is also possible and contemplated to use compounds of this invention in combination with other drugs or agents for treatment of various weight and feeding-related disorders. Compounds of this invention may be employed for decreasing food intake and/or body weight in combination with any other agent or drug heretofore employed as a diet aid, or for decreasing food intake and/or body weight. Compounds of this invention may further be employed for increasing food intake and/or body weight in combination with any other agent or drug heretofore employed for increasing food intake and/or body weight.

Drugs that reduce energy intake include, in part, various pharmacological agents, referred to as anorectic drugs, which are used as adjuncts to behavioral therapy in weight reduction programs. Classes of anorectic drugs include, but are not limited to, noradrenergic and serotonergic agents. Noradrenergic medications may be described as those medications generally preserving the anorectic effects of amphetamines but with weaker stimulant activity. The noradrenergic drugs, except phenylpropanolamine, generally act through a centrally mediated pathway in the hypothalamus that causes anorexia. Phenylpropanolamine, a racemic mixture of norephedrine esters, causes a release of norepinephrine throughout the body and stimulates hypothalamic adrenoreceptors to reduce appetite.

Suitable noradrenergic agents include, but are not limited to, diethylpropion such as TENUATE™ (1-propanone, 2-(diethylamino)-1-phenyl-, hydrochloride) commercially available from Merrell; mazindol (or 5-(p-chlorophenyl)-2,5-dihydro-3H-imidazo[2,1-a]isoindol-5-ol) such as SANOREX™ commercially available from Novartis or MAZANOR™ commercially

available from Wyeth Ayerst; phenylpropanolamine (or Benzenemethanol, alpha-(1-aminoethyl)-, hydrochloride); phentermine (or Phenol, 3-[[4,5-dihydro-1H-imidazol-2-yl)ethyl](4-methylphenyl)amino], monohydrochloride) such as ADIPEX-P™ commercially available from Lemmon, FASTIN™ commercially available from Smith-Kline Beecham and
5 Ionamin™ commercially available from Medeva; phendimetrazine (or (2S,3S)-3,4-Dimethyl-2-phenylmorpholine L-(+)-tartrate (1:1)) such as METRA™ commercially available from Forest, PLEGINE™ commercially available from Wyeth-Ayerst; PRELU-2™ commercially available from Boehringer Ingelheim, and STATOBEX™ commercially available from Lemmon; phendamine tartrate such as THEPHORIN™ (2,3,4,9-Tetrahydro-2-methyl-9-phenyl-1H-
10 indenol[2,1-c]pyridine L-(+)-tartrate (1:1)) commercially available from Hoffmann-LaRoche; methamphetamine such as DESOXYN™ Tablets ((S)-N, (alpha)-dimethylbenzeneethanamine hydrochloride) commercially available from Abbott; and phendimetrazine tartrate such as BONTRIL™ Slow-Release Capsules (-3,4-Dimethyl-2-phenylmorpholine Tartrate) commercially available from Amarin.

15 Suitable serotonergic agents include, but are not limited to, sibutramine such as MERIDIA™ capsules (a racemic mixture of the (+) and (-) enantiomers of cyclobutanemethanamine, 1-(4-chlorophenyl)-N,N-dimethyl-(alpha)-(2-methylpropyl)-, hydrochloride, monohydrate) commercially available from Knoll, fenfluramine such as Pondimin™ (Benzeneethanamine, N-ethyl-alpha-methyl-3-(trifluoromethyl)-, hydrochloride)
20 commercially available from Robbins; dexfenfluramine such as Redux™ (Benzeneethanamine, N-ethyl-alpha-methyl-3-(trifluoromethyl)-, hydrochloride) commercially available from Interneuron. Fenfluramine and dexfenfluramine stimulate release of serotonin and inhibit its reuptake. Sibutramine inhibits the reuptake of serotonin, norepinephrine and dopamine, but does not stimulate secretion of serotonin.

25 Other serotonergic agents useful with the practice of the present invention include, but are not limited to, certain auoretic gene 5HT1a inhibitors (brain, serotonin) such as carbidopa and benserazide as disclosed by U.S. Pat. No. 6,207,699 which is incorporated herein by reference; and certain neurokinin 1 receptor antagonist and selective serotonin reuptake inhibitors including fluoxetine, fluvoxamine, paroxetine, sertraline and other useful compounds

as disclosed by U.S. Pat. No. 6,162,805 which is incorporated herein by reference. Other potential agents that may be employed include, for example, 5HT_{2c} agonists.

Other useful compounds for reducing energy intake include, but are not limited to, certain aryl-substituted cyclobutylalkylamines as disclosed by U.S. Pat. No. 6,127,424 which is incorporated herein by reference; certain trifluoromethylthiophenylethylamine derivatives as disclosed by U.S. Pat. No. 4,148,923 which is incorporated herein by reference; certain compounds as disclosed by U.S. Pat. No. 6,207,699 which is incorporated herein by reference; certain kainite or AMPA receptor antagonists as disclosed by U.S. Pat. No. 6,191,117 which is incorporated herein by reference; certain neuropeptide receptor subtype 5 as disclosed by U.S. Pat. No. 6,140,354 which is incorporated herein by reference; and certain alpha-blocking agents as disclosed by U.S. Pat. No. 4,239,763 which is incorporated herein by reference.

Moreover, several peptides and hormones regulate feeding behavior. For example, cholecystokinin and serotonin act to decrease appetite and food intake. Leptin, a hormone produced by fat cells, controls food intake and energy expenditure. In obese persons who are losing weight without medications, a decrease in weight is associated with a decrease in circulating levels of leptin, suggesting its role in weight homeostasis. Obese patients with high leptin levels are thought to have peripheral leptin resistance secondary to the down-regulation of leptin receptors. Non-limiting examples of useful compounds affecting feeding behavior include certain leptin-lipolysis stimulated receptors as disclosed by WO 01/21647 which is incorporated herein by reference; certain phosphodiesterase enzyme inhibitors as disclosed by WO 01/35970 which is incorporated herein by reference; certain compounds having nucleotide sequences of the mahogany gene as disclosed by WO 00/05373 which is incorporated herein by reference; and certain sapogenin compounds as disclosed by U.S. Pat. No. 4,680,289 which is incorporated herein by reference.

Other useful compounds include certain gamma peroxisome proliferator activated receptor (PPAR) agonists as disclosed by WO 01/30343 and U.S. Pat. No. 6,033,656 which are incorporated herein by reference and certain polypeptides such as fibroblast growth factor-10 polypeptides as disclosed by WO 01/18210 which is incorporated herein by

reference.

Moreover, monoamine oxidase inhibitors that decrease energy intake or increase energy expenditure are useful with the practice of the present invention. Suitable, but non-limiting, examples of monoamine oxidase inhibitors include befloxatone, moclobemide,
5 brofaromine, phenoxathine, esuprone, befol, toloxatone, pirlindol, amiflamine, serclorephine, bazinaprine, lazabemide, milacemide, caroxazone and other certain compounds as disclosed by WO 01/12176 which is incorporated herein by reference.

Certain compounds that increase lipid metabolism are also useful with the practice of the present invention. Such compounds include, but are not limited to, evodiamine
10 compounds as disclosed by U.S. Pat. No. 6,214,831 which is incorporated herein by reference.

Nutrient partitioning agents and digestive inhibitors are another strategy in the treatment of obesity by interfering with the breakdown, digestion or absorption of dietary fat in the gastrointestinal tract. Gastric and pancreatic lipases aid in the digestion of dietary
15 triglycerides by forming them into free fatty acids that are then absorbed in the small intestine. Inhibition of these enzymes leads to inhibition of the digestion of dietary triglycerides. Non-limiting examples include a lipase inhibitor, orlistat, such as XENICAL™ capsules ((S)-2-formylamino-4-methyl-pentanoic acid (S)-1-[[[(2S, 3S)-3-hexyl-4-oxo-2-oxetanyl]methyl]-dodecyl ester) commercially available from Roche Laboratories and certain benzoxazinone
20 compounds as described by WO 00/40247 which is incorporated herein by reference.

Agents that increase energy expenditure are also referred to as thermogenic medications. Non-limiting examples of suitable thermogenic medications include xanthines, such as caffeine and theophylline, selective β -3-adrenergic agonists, for example certain compounds in U.S. Pat. No. 4,626,549 which is incorporated by reference herein, and α -2-
25 adrenergic and growth hormones compounds as described in U.S. Pat. Nos. 4,937,267 and 5,120,713 which are incorporated by reference herein.

Generally, a total dosage of the above-described obesity control agents or medications, when used in combination with a compound of this invention can range from 0.1 to 3,000 mg/day, preferably from about 1 to 1,000 mg/day and more preferably from about 1

to 200 mg/day in single or 2-4 divided doses. The exact dose, however, is determined by the attending clinician and is dependent on such factors as the potency of the compound administered, the age, weight, condition and response of the patient.

Agents or drugs employed for increasing food intake and/or body weight include
5 appetite stimulants such as megestrol acetate, adrenocorticoids such as prednisolone and dexamethasone, cyproheptidine, serotonergic drugs such as fenfluramine, neuropeptide Y, and androgen antagonists such as flutamide, nilutamide, and zanolone.

Assays and Animal Models.

Selected compounds were tested in assays to determine binding and functional
10 status, and were tested in animal models of feeding behavior as discussed below. The following assays and animal models were employed, with modifications as discussed in the examples.

Competitive inhibition assay using [125 I]-NDP- α -MSH. A competitive inhibition binding assay was performed using membrane homogenates prepared from HEK-293 cells
15 that express recombinant hMC4-R, hMC3-R, or hMC5-R, and from B-16 mouse melanoma cells (containing endogenous MC1-R). In some instances, HEK-293 cells that express recombinant hMC1-R were employed. In the examples that follow, all MC3-R, MC4-R and MC5-R values are for human recombinant receptors. MC1-R values are for B-16 mouse melanoma cells, unless the heading is "hMC1-R", in which case the value is for human
20 recombinant MC1-R. Assays were performed in 96 well GF/B Millipore multiscreen filtration plates (MAFB NOB10) pre-coated with 0.5% bovine serum albumin (Fraction V). Membrane homogenates were incubated with 0.2nM (for hMC4-R) 0.4 nM (for MC3-R and MC5-R) or 0.1 nM (for mouse B16 MC1-R or hMC1-R) [125 I]-NDP- α -MSH (Perkin Elmer) and increasing concentrations of test compounds in buffer containing 25 mM HEPES buffer (pH 7.5) with 100
25 mM NaCl, 2 mM CaCl_2 , 2 mM MgCl_2 , 0.3 mM 1,10-phenanthroline, and 0.2% bovine serum albumin. After incubation for 60 minutes at 37° C, the assay mixture was filtered and the membranes washed three times with ice-cold buffer. Filters were dried and counted in a gamma counter for bound radioactivity. Non-specific binding was measured by inhibition of binding of [125 I]-NDP- α -MSH in the presence of 1 μ M NDP- α -MSH. Maximal specific binding

(100%) was defined as the difference in radioactivity (cpm) bound to cell membranes in the absence and presence of 1 μ M NDP- α -MSH. Radioactivity (cpm) obtained in the presence of test compounds was normalized with respect to 100% specific binding to determine the percent inhibition of [125]-NDP- α -MSH binding. Each assay was conducted in triplicate and the actual mean values are described, with results less than 0% reported as 0%. Ki values for test compounds were determined using Graph-Pad Prism® curve-fitting software.

Competitive binding assay using [125]-AgRP (83-132). Competitive binding studies using [125]-AgRP (83-132) were carried out using membrane homogenates isolated from cells that express hMC4-R. The assays were performed in 96-well GF/B Millipore multiscreen filtration plates (MAFB NOB10) pre-coated with 0.5% bovine serum albumin (Fraction V). The assay mixture contained 25 mM HEPES buffer (pH 7.5) with 100 mM NaCl, 2 mM CaCl_2 , 2 mM MgCl_2 , 0.3 mM 1,10-phenanthroline, 0.5% bovine serum albumin, membrane homogenates, radioligand [125]-AgRP (83-132) (Perkin Elmer) and increasing concentrations of compounds in a total volume of 200 μ L. Binding was measured at radioligand concentrations of 0.2 nM. After incubating for 1 hour at 37° C, the reaction mixture was filtered and washed with assay buffer containing 500 mM NaCl. The dried discs were punched out from the plate and counted on a gamma counter. The total binding of the radioligand did not exceed 10% of the counts added to the reaction mixture. Ki values for test compounds were determined using Graph-Pad Prism® curve-fitting software.

Assay for agonist activity. Accumulation of intracellular cAMP was examined as a measure of the ability of the test compounds to elicit a functional response in HEK-293 cells that express MC4-R. Confluent HEK-293 cells that express recombinant hMC4-R were detached from culture plates by incubation in enzyme-free cell dissociation buffer. Dispersed cells were suspended in Earle's Balanced Salt Solution containing 10 mM HEPES (pH 7.5), 1 mM MgCl_2 , 1mM glutamine, 0.5% albumin and 0.3 mM 3-isobutyl-1-methyl-xanthine (IBMX), a phosphodiesterase inhibitor. The cells were plated in 96-well plates at a density of 0.5×10^5 cells per well and pre-incubated for 30 minutes. Cells were exposed for 1 hour at 37° C to test compounds dissolved in DMSO (final DMSO concentration of 1%) at a concentration range of 0.05 - 5000 nM in a total assay volume of 200 μ L. NDP- α -MSH was used as the

reference agonist. At the end of the incubation period, cells were disrupted by the addition of 50 μ L of lysis buffer (cAMP EIA kit, Amersham) followed by vigorous pipetting. Levels of cAMP in the lysates were determined using a cAMP EIA kit (Amersham). Data analysis was performed by nonlinear regression analysis with Graph-Pad Prism[®] software. The maximum
5 efficacies of the test compounds were compared to that achieved by the reference melanocortin agonist NDP- α MSH.

Food intake after IN and IP dosing. Changes in food intake were evaluated for selected compounds. Male C57BL/6 mice were obtained from Jackson labs (Bar Harbor, ME). Animals were individually housed in conventional plexiglass hanging cages and
10 maintained on a controlled 12 hour on/off light cycle. Water and pelleted (Harlan Teklad 2018 18% Protein Rodent Diet) food was provided *ad libitum*. The mice were dosed IP (by intraperitoneal injection) after a 24 hour fast or IN (by intranasal administration) with vehicle or selected compounds (0.1–3 mg/kg, and in some cases up to 10 mg/kg). All animals were dosed once a day (or up to four consecutive days) at the start of the "lights off" period. The
15 changes in food intake weight for the 4 hour and 20 hour period after dosing relative to control animals administered vehicle were determined.

Determination of mass and nuclear magnetic resonance analysis. The mass values were determined using a Waters MicroMass ZQ device utilizing a positive mode. Mass determinations were compared with calculated values and expressed in the form of
20 mass weight plus one (M+1 or M+H).


Proton NMR data was obtained using a Bruker 300 MHz spectrometer. The spectra were obtained after dissolving compounds in a deuteriated solvent such as chloroform, DMSO, or methanol as appropriate.

Synthetic Methods of the Invention.

25 One general strategy includes developing a linear intermediate using chiral building blocks such as amino acid derivatives. The linear intermediate can be cyclized using a Mitsunobo reaction strategy or by spontaneous cyclization through reactive groups such as a reaction between an amine and an ester or between an amine and an aldehyde function. In these cyclizations, the driving force for intramolecular reaction versus intermolecular reaction

is the thermodynamically favored reaction forming a six-membered ring structure. In many instances, the methodology incorporates conditions that do not involve inversion or racemization of chiral centers. In some instances where a small percentage of racemate is observed, such as in use of an α -amino aldehyde in certain positions, the desired chiral product is easily purified by methods known in the art, such as flash chromatography on a silica gel column.

The group containing the Q ring is preferably made by use of an aldehyde derivative of a D-amino acid. By use of an α -amino aldehyde the resulting group has, in its most basic form, the general structure:



By way of example, where an aldehyde derivative of D-Phe is employed in synthesis, in the resulting compound r is 1, and Q is phenyl. However, it can readily be seen that any D-amino acid may be employed as an aldehyde derivative, and may further be seen that rather than NH_2 any amine capping group may be employed in lieu of one or both hydrogen atoms. In synthesis, preferably an N-protected D-amino acid aldehyde is employed, where the N-protecting group is conventionally Boc or Fmoc. Because of the inherent instability of α -amino aldehydes in solution, these compounds are preferably synthesized immediately prior to use. Two different methods may be used for synthesis.

In the first method, to an N-protected amino acid (such as with a Boc- or Fmoc-group) in dichloromethane was added TBTU (1 eq.) (here and elsewhere "eq." is an abbreviation for equivalent or equivalents, as the context requires) and NMM (1 eq.). The mixture was stirred for half an hour and N,O-dimethylhydroxylamine hydrochloride (1 eq.) and NMM (1 eq.) were added. The reaction was carried out overnight. The solvent was removed and EtOAc was added. The organic phase was washed by aqueous sodium bicarbonate, brine and dried over sodium sulfate. After evaporation of solvent and drying under vacuum the residue was dissolved in THF under nitrogen at -78°C . To this solution was added LAH (1 M in THF, 1.5 eq.) slowly. The solution was stirred for an additional half hour. The reaction was diluted by ether and quenched by aqueous potassium hydrogen sulfate. The organic phase was


washed with 1 N HCl, water, brine and dried over sodium sulfate. After removal of solvent the aldehyde was used immediately for the next step reaction without purification.

In the second method, to an N-protected amino acid (such as with a Boc- or Fmoc-group) in THF was added borane-THF (1 M, 1.2 eq.) slowly at 0° C. The temperature was
5 raised to room temperature and the solution stirred for 2 hours. The reaction was quenched by 1 N HCl and the solvent was evaporated. The crude product was purified on a silica gel column to give a pure N-protected amino alcohol. This alcohol was dissolved in dry dichloromethane and Dess-Martin periodinane (1.1 eq.) was added. The solution was stirred for 1 hour and the reaction was diluted by ether. The organic phase was washed by
10 saturated sodium bicarbonate with 10% sodium thiosulfate, then water, then brine and dried over sodium sulfate. After removal of solvent the crude product was used for the next step reaction immediately without further purification.



In the synthetic methods employed, either of the foregoing methods may be employed to utilize a D-amino acid aldehyde.

15 In general, the synthetic methods employed were modifications of those described in applications cited above, including specifically Patent Application Serial No. 10/837,519, but employing an amino acid aldehyde, and in most instances a D-amino acid aldehyde.

Scheme 1



- R_2 in Fmoc-NHCH₂(R₂)-COOH is the group in position R_{2a} or R_{2b} of Structure I. Thus R₂ may be cyclohexane, methyl-cyclohexane, ethyl-cyclohexane, propyl-cyclohexane, benzene, toluene, ethyl-benzene, propyl-benzene, and the like, including specifically any

group of the general structure  or , where y is a value from 0 to 5.

To a solution of Fmoc-NHCH₂(R₂)-COOH (**1-1**) in DCM was added TBTU (1.05 eq.) and NMM (1.05 eq.), and the mixture stirred for one hour at room temperature. To this mixture were added N,O-dimethylhydroxyamine HCl salt (1.1 eq.) and NMM (1.1 eq.). The reaction was continued at room temperature overnight. The solvent was removed, and the residue partitioned between EtOAc and water. The organic layer was washed by water, 1 N HCl, saturated sodium bicarbonate, brine and dried over sodium sulfate. The solvent was removed and the crude compound **1-2** was used for next step reaction.

To a solution of compound **1-2** in dry THF was added LAH (1.2 eq.) slowly under nitrogen at -78° C. After the addition was completed the reaction mixture was stirred at -78° C for one hour. The reaction was quenched by the addition of aqueous potassium hydrogen sulfate solution. The mixture was diluted with EtOAc and the solid was removed and solvent evaporated. The residue was dissolved in EtOAc and the organic layer was washed with 1 N HCl, water and dried over sodium sulfate. The solvent was removed and the crude product **1-3** was used for the next step reaction.

To a suspension of H-Orn(Boc)-OMe HCl salt in THF was added TEA (1 eq.). The mixture was stirred for 30 minutes under nitrogen. To this mixture was added **1-3** in THF, then 4 Å molecular sieves. The mixture was stirred for 2 hours at room temperature and sodium triacetoxy borohydride (1.5 eq.) was added. The reaction was continued at room temperature overnight. The solids were removed by passing through a Celite pad, removing the solvent and partitioning the residue between EtOAc and water. The organic layer was collected and dried over sodium sulfate. After removal of the solvent the product **1-4** was obtained as a crude compound, which was used for the next step reaction without further purification.

Compound **1-4** was dissolved in 30% diethyl amine in EtOAc. The reaction was continued overnight at room temperature. The solvent was removed and the residue was purified on a silica gel column to give pure product **1-5**.

Compound **1-5** was dissolved in DCM and TEA (1.5 eq.) added. To this solution was added benzyl chloroformate (1.2 eq.) at 0° C. The reaction mixture was stirred overnight at room temperature. The solvent was removed from the reaction mixture and the residue was

purified on a silica gel column to give compound **1-6**.

To a solution of compound **1-6** THF was added borane-THF (6 eq.) slowly at 0° C. The reaction was continued at room temperature overnight. The reaction was quenched by the addition of water, and the solvent then removed. The residue was stirred in sodium
5 hydroxide (10 eq.) in methanol for 24 hours, and EtOAc and water then added. The organic layer was washed with water and brine and dried over sodium sulfate. After removal of the solvent the residue was purified on a silica gel column to give pure product **1-7**.

Formation of compound **1-8**:

Method A: A solution of J-carboxylic acid (1.5 eq.) HOAt (1.5 equiv) and EDC (1.5
10 eq.) in N, N-dimethylformamide was stirred at 0° C for 30 minutes. To this solution was added compound **1-7**. The reaction was continued overnight at room temperature. The solvent was removed and the residue purified by flash chromatography to give compound **1-8**.


Method B: To a solution of compound **1-7** and TEA (3 eq.) in THF was added J-carbonyl chloride (1.5 eq.) at 0° C. The reaction was continued at room temperature
15 overnight. The solvent was removed and the residue purified by silica gel column to give product **1-8**.

Compound **1-8** was treated with a solution of TFA/DCM (v:v = 3:1) for one hour. The solvent was removed and the residue was dissolved in acetonitrile. This solution was basified by addition of TEA. The solvent was removed and residue re-dissolved in acetonitrile. To
20 this solution was added TEA (1 eq.) and N,N'-bis(t-butoxycarbonyl)-1H-pyrazole-1-carboxamide (1.2 eq.), and the resulting mixture stirred at room temperature overnight. After removal of the solvent the residue was purified on a silica gel column to give product **1-9**.



Compound **1-9** was dissolved in ethanol and stirred at one atmosphere of hydrogen
25 in the presence of a catalytic amount of Pd/C (10%). The reaction was continued at room temperature overnight. The catalyst was removed by filtration. The solvent was removed to give crude product. This crude product was dissolved in THF and reacted with Q-aldehyde (Q-L₃-) derived from Q-COOH in the manner described for the formation of compound **1-4**. Purification by silica gel column gave a Boc-protected compound, which was subjected to

treatment with TFA/DCM (50:50) for one hour. After evaporation of solvent the final compound **1-10** was purified by HPLC.

Scheme 2



- 5 R₂ in Fmoc-NHCH₂(R₂)-COOH (**2-1**) is the group in position R_{2a} or R_{2b} of Structure I. Thus R₂ may be cyclohexane, methyl-cyclohexane, ethyl-cyclohexane, propyl-cyclohexane, benzene, toluene, ethyl-benzene, propyl-benzene, and the like, including specifically any

group of the general structure  or , where y is a value from 0 to 5.

To a solution of Fmoc-NHCH₂(R₂)-COOH (**2-1**) in DCM were added TBTU (1.05 eq.) and NMM (1.05 eq.). The mixture was stirred for one hour at room temperature. To this mixture were added N,O-dimethylhydroxylamine HCl salt (1.1 eq.) and NMM (1.1 eq.). The reaction was continued at room temperature overnight. The solvent was removed. The residue was partitioned between EtOAc and water. The organic layer was washed by water, 1 N HCl, saturated sodium bicarbonate, brine and dried over sodium sulfate. The solvent was removed and the crude compound **2-2** was used for next step reaction.

To a solution of compound **2-2** in dry THF was added LAH (1.2 eq.) slowly under nitrogen at -78° C. After completion of this addition, the reaction mixture was stirred at -78° C for one hour. The reaction was quenched by addition of aqueous potassium hydrogen sulfate solution. The mixture was diluted with EtOAc and the solid was removed. The solvent was evaporated and the residue was dissolved in EtOAc and the organic layer was washed with 1 N HCl, water and dried over sodium sulfate. The solvent was removed and the crude product **2-3** was used for next step reaction.

To a suspension of H-Orn(Boc)-OMe HCl salt in THF was added TEA (1 eq.). The mixture was stirred for 30 minutes under nitrogen. To this mixture was added **2-3** in THF, and then 4 Å molecular sieves. The mixture was stirred for 2 hours at room temperature and sodium triacetoxy borohydride (1.5 eq.) was then added. The reaction was continued at room temperature overnight. The solids were removed by passing through a Celite pad. The solvent was removed and the residue was partitioned between EtOAc and water. The organic layer was collected and dried over sodium sulfate. After removal of solvent the product **2-4** was obtained as a crude compound, which was used for next step reaction without further purification.

Compound **2-4** was dissolved in 30% diethyl amine in EtOAc. The reaction was continued overnight at room temperature. The solvent was removed and the residue was purified on a silica gel column to give pure product **2-5**.

Compound **2-5** was dissolved in DCM and TEA (1.5 eq.) was added. To this solution was added benzyl chloroformate (1.2 eq.) at 0° C. The reaction mixture was stirred overnight at room temperature. The solvent was removed from the reaction mixture and the residue was purified on a silica gel column to give compound **2-6**.

5 To a solution of compound **2-6** THF was added borane-THF (6 eq.) slowly at 0° C. The reaction was continued at room temperature overnight. The reaction was quenched by addition of water, followed by solvent removal. The residue was stirred in sodium hydroxide (10 eq.) in methanol for 24 hours. EtOAc and water were added. The organic layer was washed with water, brine and dried over sodium sulfate. After removal of the solvent the
10 residue was purified on a silica gel column to give pure product **2-7**.

Formation of compound **2-8**:

Method A: A solution of J-carboxylic acid (1.5 eq.) HOAt (1.5 eq.) and EDC (1.5 eq.) in N, N-dimethylformamide was stirred at 0° C for 30 minutes. To this solution was added compound **2-7**. The reaction was continued overnight at room temperature. The solvent was
15 removed and residue purified by flash chromatograph to give compound **2-8**.


Method B: To a solution of compound **2-7** and TEA (3 eq.) in THF was added J-carbonyl chloride (1.5 eq.) at 0° C. The reaction was continued at room temperature overnight. The solvent was removed and the residue purified by silica gel column to give product **2-8**.


20 Compound **2-8** was treated with a solution of TFA/DCM (v:v = 3:1) for one hour. The solvent was removed and the residue dissolved in acetonitrile. This solution was basified by addition of TEA. The solvent was removed and residue was re-dissolved in acetonitrile. To this solution was added TEA (1 eq.) and N,N'-bis(t-butoxycarbonyl)-1H-pyrazole-1-carboxamidine 1.2 eq.). The mixture was stirred at room temperature overnight. After
25 removal of solvent the residue was purified by a silica gel column to give product **2-9**.

Compound **2-9** was dissolved in ethanol and stirred at one atmosphere of hydrogen in the presence of a catalytic amount of Pd/C (10%). The reaction was continued at room temperature overnight, and the catalyst removed by filtration. The solvent was removed to give a crude product. This crude product was dissolved in THF and reacted with Q-COOH in

the manner described in the formation of compound **2-8** (Method A). Purification by silica gel column gave a Boc-protected compound, which was subjected to treatment with TFA/DCM (50:50) for one hour. After evaporation of solvent the final compound **2-10** was purified by HPLC.

5 **Scheme 3**





A mixture of Cbz-Glu(OtBu)-OH, TBTU (1.1 eq.) and NMM (1.5 eq.) in 100 mL of DCM were stirred at room temperature under nitrogen for 30 minutes. To the solution were added $\text{NH}_2\text{-CH(R}_2\text{)-COOMe}$ in hydrochloride salt (1.05 eq.) and NMM (1.13 eq.). The mixture was stirred at room temperature overnight. The solvent was removed and the residue was dissolved in 250 mL EtOAc. The organic solvent was washed with water, 1 N HCl, saturated aqueous sodium bicarbonate solution, water and dried over sodium sulfate. After removal of solvent, the product (**3-1**) was used for the next step reaction without further purification.

Compound **3-1** was dissolved in EtOAc. It was subject to the treatment with hydrogen at 1 atmosphere in the presence of Pd/C at room temperature for three days. The reaction mixture was filtered through a Celite pad, which was then washed with methanol. The solvent removed and the product **3-2** was used for the next step reaction without purification.

Compound **3-2** was dissolved in DMF. The solution was heated at 90° C for three days. The solvent was removed and the residue was dissolved in DCM, which was washed with 1 N HCl. The organic layer was separated and dried over sodium sulfate. After solvent

removal, product **3-3** was obtained.

A suspension of compound **3-3** in THF was stirred at 0° C. To the suspension was added LAH (4.6 eq.). The mixture was stirred for 25 minutes at 0° C and 4 hours at room temperature and then refluxed under nitrogen overnight. The reaction was quenched by the addition water, 15% sodium hydroxide and water at 0° C. The mixture was stirred at room temperature for an additional 30 minutes. The solid was removed by filtration and washed with ether. The solvent was removed and dried under vacuum to give the crude compound **3-4**.

Compound **3-4** was dissolved in THF. To this solution was added benzyl chloroformate (2.5 eq.) followed by water and sodium bicarbonate (6 eq.), such that the ratio of THF to water was 2 to 1. The mixture was stirred overnight at room temperature. To this mixture were added EtOAc and water. The organic layer was washed with water and dried over sodium sulfate. The solvent was removed and the residue was dissolved in methanol and 1 N sodium hydroxide (3 eq.). The reaction was carried out at room temperature for three days. The solvent was removed and the resulted residue was dissolved in EtOAc. The organic phase was washed with 1 N HCl, water, aqueous sodium bicarbonate and dried over sodium sulfate. The solvent was removed and the residue purified on a silica gel column to give product **3-5**.

Compound **3-5** was dissolved in DCM. To this solution was added Dess-Martin periodinane (1.1 eq.). The reaction was continued at room temperature for 1.5 hours. Ether was added to dilute the mixture and the reaction was quenched by addition of a solution of sodium thiosulfate in saturated sodium bicarbonate. The organic layer was washed with the same solution and dried over sodium sulfate. After removal of solvent the crude product **3-6** was used for next step reaction without further purification.

To a solution of compound **3-6** and N-Boc-ethylenediamine (1.05 eq.) in THF was added molecular sieves. The mixture was stirred for 3 hours at room temperature. To this mixture was added sodium triacetoxyborohydride (1.5 eq.). The reaction mixture was stirred at room temperature overnight, and the solid removed by filtration. After the solvent was removed the residue was partitioned between EtOAc and water, the organic layer separated


and the water layer extracted with EtOAc. The combined organic layer was dried over sodium sulfate. After removal of solvent the residue was dissolved in THF and water (v:v = 2:1). To this solution were added di-t-butyl dicarbonate (1.2 eq.) and sodium bicarbonate (5 eq.). The mixture was stirred overnight at room temperature. EtOAc was added and the
5 organic layer was washed with water and dried over sodium sulfate. After removal of solvent the residue was purified on silica gel column to give product **3-7**.


Compound **3-7** was dissolved in ethanol and treated with hydrogen under atmosphere pressure in the presence of catalytic amount of Pd/C. The reaction was carried out overnight at room temperature. The solid was filtered and washed with ethanol several
10 times. The solvent was removed to give product **3-8**.

To a solution of Q-COOH (4 eq.) in DMF was added HOAT (4 eq.) and EDC (4 eq.) at 0° C. After the mixture was stirred for 30 minutes, compound **3-8** was added to this mixture. The reaction was continued overnight at room temperature. The solvent was removed and the residue was purified on a silica gel column to give product **3-9**.

15 Compound **3-9** was treated by TFA for three hours at room temperature. After removal of solvent, the residue was purified by HPLC to give compound **3-10**.

Scheme 4





Fmoc-Lys(Trt)-OH was partially dissolved in DCM. To this mixture was added TBTU (1.1 eq.) and NMM (1.5 eq.). After this mixture was stirred at room temperature under

nitrogen for 45 minute, and $\text{NH}_2\text{-CH(R}_2\text{)-COOMe}$ (i.e. H-D-Leu-OMe.HCl) (1.05 eq.) and NMM (1.1 eq.) were added. The reaction was carried out at room temperature overnight.

The solvent was evaporated and the residue was partitioned between EtOAc and water. The organic layer was washed by 1 N HCl, saturated NaHCO_3 , water and dried over sodium sulfate. After removal of solvent, product 1.1 was obtained and was used for next step reaction without further purification.

Compound **4-1** was dissolved in 30% Et_2NH in EtOAc. The solution was stirred at room temperature for 2 hours. The solvent was removed. The crude product **4-2** was used for next step reaction without further purification.

Compound **4-2** was partially dissolved in dry DMF. The mixture was heated at 90°C under nitrogen for three days. The reaction was stopped and DMF was removed under vacuum. This crude product was purified on silica gel column eluted with EtOAc/heptane (1:1), DCM, and subsequently MeOH/DCM (9:1). The final product **4-3** was collected after evaporation of solvent.

Compound **4-3** was suspended in THF at 0°C under nitrogen. To this suspension was added LAH (3.5 eq.) dropwise. The suspension became a clear solution after addition of LAH was completed. The reaction was stirred at room temperature for 45 minutes and refluxed overnight. The reaction mixture was quenched by addition of water, 15% NaOH and water, in sequence at 0°C . The mixture was stirred for 20 minutes at room temperature. The solid was removed by filtration and washed with ether, with ether evaporated to give the crude product **4-4**.

Compound **4-4** was dissolved in THF. To this solution benzyl chloroformate (3 eq.) was added, and followed by water and sodium bicarbonate (5 eq.). The reaction mixture was stirred at room temperature for 3 hours. To this mixture water and EtOAc was added. The organic layer was separated and washed by water till the aqueous layer reached neutral pH. The organic layer was dried over sodium sulfate. The solvent was evaporated and the residue was purified on silica gel column eluted with EtOAc/heptane (1:4) to give product **4-5**.

Compounds **4-5** was dissolved in a solution of 5% TFA/1% TIS/DCM, and the mixture stirred for 1 hour. The reaction mixture was diluted with DCM. The organic phase was

washed with saturated sodium bicarbonate, water, brine and dried over sodium sulfate. After evaporation of solvent the crude product **4-6** was used for next step reaction.

Compound **4-6** was dissolved in DCM. To this solution was added pyridine (10 eq.), and then 2-nitrobenzenesulfonyl chloride (2 eq.) slowly at 0° C. The temperature was raised
5 to room temperature and stirred overnight. The solvent was removed. The residue was dissolved in EtOAc, which was washed by 1N HCl, water, brine and dried over sodium sulfate. After removal of solvent, the residue was purified on silica gel column eluted with 25% EtOAc in heptane. After removal of solvent, compound **4-7** was obtained.

Compound **4-7**, TPP (3 eq.) and N-Boc-2-hydroxy-ethylamine (3 eq.) were dissolved
10 in dry toluene. To this solution was added DIAD (3 eq.) in toluene at 0° C. The temperature was raised to room temperature after 30 minutes and the solution was stirred overnight. The solvent was removed and the residue purified on silica gel column eluted with 50% EtOAc in heptane. Compound **4-8** was obtained after removal of solvent.

Compound **4-8** was dissolved in dry acetonitrile. To this solution was added
15 potassium carbonate (6 eq.) and 4-mercaptophenol (4.5 eq.). The mixture was stirred overnight at room temperature. The solvent was removed, and the residue partitioned with EtOAc and water. The organic layer was separated and washed by water, brine and dried over sodium sulfate. The crude compound **4-9** was used for next step reaction.

Compound **4-9** was dissolved in THF/water (2/1). To this solution sodium
20 bicarbonate (5 eq.) and di-t-butyl carbonate (2 eq.) were added subsequently. The mixture was stirred overnight at room temperature. After removal of THF, EtOAc was added to extract the product. The organic layer was washed with 1 N HCl, water, brine and dried over sodium sulfate. The solvent was removed and the residue purified on silica gel column eluted with 25% EtOAc in heptane. Product **4-10** was obtained after removal of solvent.

25 Compound **4-10** was treated with catalytic amounts of Pd on carbon in ethanol under hydrogen (1 atmosphere) overnight at room temperature. After filtration through a Celite pad, the solvent was removed. The crude product was dried under vacuum and it was used for next step reaction without further purification.

Q-COOH (i.e. Boc-D-2-Naphthylalanine) (4 eq.), EDC (4 eq.) and HOAt (4eq.) were


dissolved in DMF. The mixture was stirred at 0° C for 30 minutes. To this solution was added Compound **4-10** in one portion. The reaction was carried out overnight at room temperature. The solvent was removed and the residue was purified on silica gel column to give product **4-11**.

5 Compound **4-11** was treated with TFA for 3 hours at room temperature. After removal of solvent the residue was purified by HPLC to give product **4-12**.

Representative Compounds of the Invention

10 **Example 1** N-{3-[(2S,5R)-1-((R)-2-Amino-3-naphthalen-2-yl-propionyl)-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe as NHCH₂(R₂)-COOH, and Boc-D-2'-naphthylalanine as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 641 (M+H).



15

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
4%	57%	100%	84%

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	250	3	171

Ki (nM) (AgRP)	
MC3-R	MC4-R
>1000	ND


In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 87% of the maximum achieved with NDP- α -MSH.

In mouse model IP feeding studies at 3 and 10 mg/kg dose levels, an 8% and 23% decrease, respectively, was observed in food intake for a period of 20 hours. In mouse model IN feeding studies at a 3 mg/kg dose level, a 15% decrease was observed in food intake for a period of 20 hours.

Example 2 **N-{3-[(2S,5R)-5-Benzyl-1-((R)-2-dimethylamino-3-naphthalen-2-yl-propionyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine**

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe as NHCH₂(R₂)-COOH, and Fmoc-D-2'-naphthylalanine as Q-COOH. Methylation of the amine proceeded by dissolving compound **2-9** in ethanol and stirring at one atmosphere of hydrogen in the presence of a catalytic amount of Pd/C (10%). The reaction was continued at room temperature overnight, and the catalyst removed by filtration. The solvent was removed to give a crude product. This crude product was dissolved in THF and reacted with Q-COOH in the manner described in the formation of compound **2-8** (Method A). Purification by silica gel column gave a Fmoc- and Boc-protected compound. The Fmoc was removed by 30% diethylamine in EtOAc for two hours. The solvent was removed and the residue was dissolved in dichloroethane. To this solution was added formaldehyde (37% aq. solution, 10 eq.). After 10 minutes stirring, sodium triacetoxyborohydride (5 eq.) was added. The mixture was then stirred overnight at room temperature. The reaction was washed with water, brine and dried over sodium sulfate. After solvent removal, the product was subjected to treatment with TFA/DCM (50:50) for one hour, and the product was purified by HPLC to give the purified compound. Following purification, the compound was tested as described above with the results shown. The mass

was analyzed as 669.3 (M+H).



Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
17	58	97	54

Ki (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	590	26	678


Ki (nM) (AgRP)

MC3-R	MC4-R
>1000	49

5 In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 90% of the maximum achieved with NDP- α -MSH.

Example 3 N-{3-[(2S,5R)-5-Benzyl-1-(4'-chloro-biphenyl-2-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

10 The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and 4'-chloro-biphenyl-2-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 658.2 (M+H).



Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
32	21	88	4

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
317	>1000	32	910


K_i (nM) (AgRP)

MC3-R	MC4-R
>1000	180

In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 106% of the maximum achieved with NDP- α -MSH.

Example 4 N-{3-[(2S,5R)-5-Benzyl-1-(4'-chloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and 4'-chloro-biphenyl-3-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 658.3 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
37	23	88	18

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
338	>500	30	>1000

K_i (nM) (AgRP)


MC3-R	MC4-R
>1000	152

In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 98% of the maximum achieved with NDP- α -MSH.

Example 5 N-{3-[(2S,5R)-5-Benzyl-1-(4'-chloro-biphenyl-4-carbonyl)-4-(2-naphthalen-


2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe as NHCH₂(R₂)-COOH, and 4'-chloro-biphenyl-4-carboxylic acid. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 658.2 (M+H).




Example 6 N-{3-[(2S,5R)-5-Benzyl-1-[2-(4'-chloro-biphenyl-2-yl)-acetyl]-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as $\text{NHCH}_2(\text{R}_2)\text{-COOH}$, and (4'-chloro-biphenyl-2-yl)-acetic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 672.2 (M+H).



Example 7 N-{3-[(2S,5R)-5-Benzyl-1-(2',4'-dichloro-biphenyl-2-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as $\text{NHCH}_2(\text{R}_2)\text{-COOH}$, and 2',4'-dichloro-biphenyl-2-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 692.3 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
0	19	90	34

 K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	>1000	54	>1000


 K_i (nM) (AgRP)

MC3-R	MC4-R
>1000	76

In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 99% of the maximum achieved with NDP- α -MSH.

Example 8 N-{3-[(2S,5R)-5-Benzyl-1-(3',4'-dichloro-biphenyl-2-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as $\text{NHCH}_2(\text{R}_2)\text{-COOH}$, and 3',4'-dichloro-biphenyl-2-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 692.3 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
3	24	94	46

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	432	23	684


K_i (nM) (AgRP)

MC3-R	MC4-R
>1000	34

In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 78% of the maximum achieved with NDP- α -MSH.

Example 9 **N-{3-[(2S,5R)-5-Benzyl-1-(3',5'-dichloro-biphenyl-2-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine**

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and 3',5'-dichloro-biphenyl-2-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 692.3 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
17	22	88	29

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	>1000	84	960


K_i (nM) (AgRP)

MC3-R	MC4-R
>1000	132

In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 86% of the maximum achieved with NDP- α -MSH.

Example 10 N-{3-[(2S,5R)-1-((R)-2-Amino-3-naphthalen-2-yl-propyl)-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 1 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and Boc-D-2'-naphthylalanine carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 627.5 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
22	49	92	87

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	>1000	61	178

K_i (nM) (AgRP)


MC3-R	MC4-R
>1000	90

In a cAMP assay for determination of functionality as to MC4-R, it was determined
 5 that the compound showed no response at a concentration of 1 μ M,

In mouse model IP feeding studies at 3 and 10 mg/kg dose levels, a 9% and 31%
 decrease, respectively, was observed in food intake for a period of 20 hours.


Example 11 N-{3-[(2S,5R)-5-Benzyl-1-(2',4'-dichloro-biphenyl-3-carbonyl)-4-(2-
 10 naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 1 using 2-
 naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and 2',4'-dichloro-biphenyl-
 3-carboxylic acid as Q-COOH, except that the guanidine Boc groups were not removed in
 step 2-9. Following purification, the compound was tested as described above with the
 15 results shown. The mass was analyzed as 892 (M+H).



Example 12 N-{3-[(2S,5R)-5-Benzyl-1-(2',4'-dichloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine


5 The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as $\text{NHCH}_2(\text{R}_2)\text{-COOH}$, and 2',4'-dichloro-biphenyl-3-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 692.3 (M+H).



10


Example 13 N-{3-[(2S,5R)-5-Benzyl-1-(3',4'-dichloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

15 The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as $\text{NHCH}_2(\text{R}_2)\text{-COOH}$, and 3',4'-dichloro-biphenyl-3-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 692.3 (M+H).



Example 14 N-{3-[(2S,5R)-5-Benzyl-1-(3',5'-dichloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine


5 The following compound was synthesized by the method of Scheme 2 using 2-naphtylacetic acid as J-COOH, D-Phe-OH as $\text{NHCH}_2(\text{R}_2)\text{-COOH}$, and 3',5'-dichloro-biphenyl-3-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 692.6 (M+H).



10


Example 15 N-{3-[(2S,5R)-5-Benzyl-1-(3-iodo-benzoyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphtylacetic acid as J-COOH, D-Phe-OH as $\text{NHCH}_2(\text{R}_2)\text{-COOH}$, and 3-iodo-phenyl-
 15 carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 674.2 (M+H).



Example 16 N-{3-[(2S,5R)-5-Benzyl-1-(2',4'-dichloro-biphenyl-3-ylmethyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

5 The following compound was synthesized by the method of Scheme 1 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and 2',4'-dichloro-biphenyl-3-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 678.6 (M+H).




Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
22%	40%	37%	8%

Example 17 N-{3-[(2S,5R)-5-Benzyl-1-(4'-chloro-biphenyl-2-ylmethyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 1 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and 4'-chloro-biphenyl-2-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 644.6 (M+H).




Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
18%	37%	46%	24%

Example 18 N-{3-[(2S,5R)-5-Benzyl-1-(4'-chloro-biphenyl-3-ylmethyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine


The following compound was synthesized by the method of Scheme 1 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and 4'-chloro-biphenyl-3-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 644.6 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
21%	35%	46%	26%

Example 19 N-{3-[(2S,5R)-5-Benzyl-1-biphenyl-2-ylmethyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine


The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as $\text{NHCH}_2(\text{R}_2)\text{-COOH}$, and biphenyl-2-carboxylic acid as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 610.5 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
0%	50%	57%	58%

Example 20 N-{3-[(2S,5R)-1-[(R)-2-Amino-3-(2,4-dimethyl-phenyl)-propyl]-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 1 using 2-naphthylacetic acid as J-COOH, Fmoc-D-Phe-OH as NHCH₂(R₂)-COOH, and Boc-D-2,4-dimethyl-D-Phe-OH as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 605 (M+H).



Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
0%	79%	96%	94%

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	78	9	115

K_i (nM) (AgRP)


MC3-R	MC4-R
876	19

In a cAMP assay for determination of functionality, it was determined that the compound showed no response at a 1 μ M concentration as to MC4-R.

In mouse model IP feeding studies at 1 and 10 mg/kg dose levels, a maximum 14% and 38% decrease, respectively, was observed in food intake for a period of 20 hours.

Example 21 N-{3-[(2S,5S)-1-((R)-2-Amino-3-naphthalen-2-yl-propionyl)-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and Boc-D-2'-naphthylalanine as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 641 (M+H).



Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
59	78	87	83


Ki (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
190	162	134	291

In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 40% of the maximum achieved with NDP- α -MSH.

Example 22 N-{3-[(2R,5R)-1-[(R)-2-Amino-3-naphthalen-2-yl-propionyl]-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 2 using 2-naphthylacetic acid as J-COOH, D-Phe-OH as NHCH₂(R₂)-COOH, and Boc-D-2'-naphthylalanine as Q-COOH, with D-Orn(Boc)-OMe used in step 2-4. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 641.3 (M+H).



Inhibition at 1 μ M (NDP- α -MSH)


MC1-R	MC3-R	MC4-R	MC5-R
19	39	80	44

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	300	121	493

Example 23 N-{3-[(2S,5R)-1-[(R)-2-Amino-3-(2,4-dimethyl-phenyl)-propyl]-5-cyclohexylmethyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 1 using 2-naphthylacetic acid as J-COOH, Fmoc-D-Cha-OH as Fmoc-NHCH₂(R₂)-COOH, and Boc-D-2,4-dimethyl-phenylalanine as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 610.8 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
12%	15%	90%	28%

 K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	494	89	269


 K_i (nM) (AgRP)

MC3-R	MC4-R
>1000	89

In a cAMP assay for determination of functionality as to MC4-R, it was determined
 5 that the compound showed no response at a concentration of 1 μ M.

Example 24 N-{3-[(2S,5R)-1-[(R)-2-Amino-3-(2,4-dichloro-phenyl)-propyl]-5-cyclohexylmethyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

10 The following compound was synthesized by the method of Scheme 1 using 2-naphthylacetic acid as J-COOH, Fmoc-D-Cha-OH as Fmoc-NHCH₂(R₂)-COOH, and Boc-D-2,4-dichloro-phenylalanine as Q-COOH. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 650.6 (M+H).

Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
14%	27%	90%	42%

K_i (nM) (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
>1000	411	100	325

K_i (nM) (AgRP)

MC3-R	MC4-R
>1000	74

In a cAMP assay for determination of functionality in a cell line that expresses MC4-R, the maximum effect observed with the compound was 22% of the maximum achieved with NDP- α -MSH.


In mouse model IP feeding studies at a 1 mg/kg dose level, an 11% decrease was observed in food intake at 20 hours.

Example 25 N-{3-[(2S,5R)-5-Cyclohexylmethyl-1-[(R)-3-(2,4-dichloro-phenyl)-2-dimethylamino-propyl]-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine

The following compound was synthesized by the method of Scheme 1 using 2-naphthylacetic acid as J-COOH, Fmoc-D-Cha-OH as Fmoc-NHCH₂(R₂)-COOH, and Fmoc-D-

2,4-dichloro-phenylalanine as Q-COOH. Methylation of the amine proceeded by dissolving compound **1-9** in ethanol and stirring at one atmosphere of hydrogen in the presence of a catalytic amount of Pd/C (10%). The reaction was continued at room temperature overnight, and the catalyst removed by filtration. The solvent was removed to give a crude product.

- 5 This crude product was dissolved in THF and reacted with Q-aldehyde derived from Q-COOH in the manner described in the formation of compound **1-4**. Purification by silica gel column gave a Fmoc- and Boc-protected compound. The Fmoc was removed by 30% diethylamine in EtOAc for two hours. The solvent was removed and the residue was dissolved in dichloroethane. To this solution was added formaldehyde (37% aq. solution, 10 eq.). After
 10 10 minutes stirring, sodium triacetoxyborohydride (5 eq.) was added. The mixture was then stirred overnight at room temperature. The reaction was washed with water, brine and dried over sodium sulfate. After solvent removal, the product was subjected to treatment with TFA/DCM (50:50) for one hour, and the product was purified by HPLC to give the purified compound. Following purification, the compound was tested as described above with the
 15 results shown. The mass was analyzed as 678.7 (M+H).



Inhibition at 1 μ M (NDP- α -MSH)

MC1-R	MC3-R	MC4-R	MC5-R
67%	75%	98%	91%

Ki (nM) (NDP- α -MSH)			
MC1-R	MC3-R	MC4-R	MC5-R
146	198	10	143

Ki (nM) (AgRP)	
MC3-R	MC4-R
241	13

In a cAMP assay for determination of functionality, it was determined that the compound showed no response at a concentration of 1 μ M.


5 In mouse model IP feeding studies at a 3 mg/kg dose level, a 3% decrease was observed in food intake at 20 hours.

Example 26 (R)-2-Amino-1-((2R,5S)-4-[(R)-2-amino-3-(2,4-dimethyl-phenyl)-propionyl]-5-[3-(2-amino-ethylamino)-propyl]-2-cyclohexylmethyl-piperazin-1-yl]-3-(2,4-dimethyl-phenyl)-propan-1-one

The following compound was synthesized by the method of Scheme 3 using Boc-2,4-dimethyl-phenylalanine as Q-COOH and D-cyclohexylalanine as $\text{NH}_2\text{CH}(\text{R}_2)\text{-COOMe}$.

Following purification, the compound was tested as described above with the results shown.

The mass was analyzed as 632.8 (M+H).



Inhibition at 1 μ M (NDP- α -MSH)

hMC1-R	MC3-R	MC4-R	MC5-R
35	22	83	43

K_i (nM) (NDP- α -MSH)


MC1-R	MC3-R	MC4-R	MC5-R
ND	ND	81	ND

Example 27 (R)-2-Amino-1-((2R,5S)-4-[(R)-2-amino-3-(2,4-dichloro-phenyl)-propionyl]-5-[3-(2-amino-ethylamino)-propyl]-2-cyclohexylmethyl-piperazin-1-yl]-3-(2,4-dichloro-phenyl)-propan-1-one

The following compound was synthesized by the method of Scheme 3 using Boc-2,4-dichloro-phenylalanine as Q-COOH and D-cyclohexylalanine as NH₂CH(R₂)-COOMe.

Following purification, the compound was tested as described above with the results shown.

The mass was analyzed as 712.6 (M+H).


Inhibition at 1 μ M (NDP- α -MSH)

hMC1-R	MC3-R	MC4-R	MC5-R
27%	36%	83%	31%

Ki (nM) (NDP- α -MSH)			
MC1-R	MC3-R	MC4-R	MC5-R
ND	ND	74	ND

Example 28 (R)-2-Amino-1-[(2R,5S)-5-[3-(2-amino-ethylamino)-propyl]-4-((R)-2-amino-3-naphthalen-2-yl-propionyl)-2-cyclohexylmethyl-piperazin-1-yl]-3-naphthalen-2-yl-propan-1-one

The following compound was synthesized by the method of Scheme 3 using Boc-D-2'-naphthylalanine acid as Q-COOH and D-cyclohexylalanine as NH₂CH(R₂)-COOMe. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 676.6 (M+H).




Ki (nM) (NDP- α -MSH)			
hMC1-R	MC3-R	MC4-R	MC5-R
22	36	81	30

Ki (nM) (NDP- α -MSH)			
MC1-R	MC3-R	MC4-R	MC5-R
ND	ND	99	ND

Example 29 (R)-2-Amino-1-[(2S,5R)-2-[4-(2-amino-ethylamino)-butyl]-4-((R)-2-amino-3-naphthalen-2-yl-propionyl)-5-cyclohexylmethyl-piperazin-1-yl]-3-naphthalen-2-yl-propan-1-one

The following compound was synthesized by the method of Scheme 4 using Boc-D-2'-naphthylalanine as Q-COOH and D-cyclohexylalanine as $\text{NH}_2\text{CH}(\text{R}_2)\text{-COOMe}$. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 691.3 (M+H).




Ki (nM) (NDP- α -MSH)

hMC1-R	MC3-R	MC4-R	MC5-R
153	435	20	103

In mouse model IN feeding studies at a 0.1 and 0.3 mg/kg dose level, a 14% and 18% decreased, respectively, in food intake was observed at 20 hours.

Example 30 (R)-2-Amino-1-[(2S,5R)-4-[(R)-2-amino-3-(2,4-dichloro-phenyl)-propionyl]-2-[4-(2-amino-ethylamino)-butyl]-5-cyclohexylmethyl-piperazin-1-yl]-3-(2,4-dichloro-phenyl)-propan-1-one

The following compound was synthesized by the method of Scheme 4 using Boc-2,4-dichloro-phenylalanine as Q-COOH and D-cyclohexylalanine as $\text{NH}_2\text{CH}(\text{R}_2)\text{-COOMe}$. Following purification, the compound was tested as described above with the results shown. The mass was analyzed as 727 (M+H).



Ki (nM) (NDP- α -MSH)

hMC1-R	MC3-R	MC4-R	MC5-R
270	443	17	139


The preceding examples can be repeated with similar success by substituting the generically or specifically described reactants and/or synthetic conditions of this invention for those used in the preceding examples.

Although the invention has been described in detail with particular reference to these preferred embodiments, other embodiments can achieve the same results. Variations and modifications of the present invention will be obvious to those skilled in the art and it is intended to cover all such modifications and equivalents. The entire disclosures of all references, applications, patents, and publications cited above, and of the corresponding application(s), are hereby incorporated by reference.

CLAIMS

What is claimed is:

1. A compound having the formula of structure I:



or an enantiomer, stereoisomer or diastereoisomer thereof, or a pharmaceutically acceptable salt thereof,

wherein

J is a ring structure selected from the group consisting of substituted or unsubstituted aromatic carbocyclic rings, substituted or unsubstituted non-aromatic carbocyclic rings, substituted or unsubstituted aromatic fused carbobicyclic ring groups, two substituted or unsubstituted aromatic carbocyclic rings wherein the rings are joined by a bond, -CH₂-, or -O-, and substituted or unsubstituted aromatic fused heterobicyclic ring groups, wherein in each instance the rings include 5 or 6 ring atoms;

W is a heteroatom unit with at least one cationic center, hydrogen bond donor or hydrogen bond acceptor wherein at least one heteroatom is nitrogen or oxygen;


Q is an aromatic carbocyclic ring selected from the group consisting of phenyl, substituted phenyl, naphthyl and substituted naphthyl;

L₁ is a bond or a linker unit comprising from one to eight backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

L₂ is a bond or -(CH₂)_z;

L₃ is a bond or a linker unit comprising from one to nine backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

R_{1a}, R_{1b}, R_{2a} and R_{2b} are selected such that:




the remaining of R_{2a} and R_{2b} is hydrogen and R_{1a} and R_{1b} are each hydrogen, R_{1a} and R_{1b} together form $=O$, or one of R_{1a} and R_{1b} is a C_1 to C_6 aliphatic linear or branched chain and the remaining of R_{1a} and R_{2b} is hydrogen,

or R_{2a} and R_{2b} are each hydrogen

5

and one of R_{1a} and R_{1b} are



or

,

and the remaining of R_{1a} and R_{1b} is hydrogen;

X is CH_2 , $C=O$ or $C=S$;


z is an index value from 1 to 6; and

y is an index value from 0 to 5;


10

wherein the carbon atom marked with an asterisk can have any stereochemical configuration.


2. The compound of claim 1 wherein J is




,




,



,




,




,


84




,




,




,



,



, or




,

unsubstituted or substituted with one or more ring substituents.

3. The compound of claim 2 wherein J is substituted with one or more ring
 5 substituents independently selected from the group consisting of hydroxyl, halogen, sulfonamide, alkyl, -O-alkyl, aryl, and -O-aryl.

4. The compound of claim 1 wherein Q is



- 10 and wherein R_{3a} , R_{3b} and R_{3c} are optional ring substituents, and when one or more are present, are the same or different and independently hydroxyl, halogen, alkyl, -O-alkyl, aryl or -O-aryl groups.


5. The compound of claim 4 wherein at least one of R_{3a} , R_{3b} or R_{3c} is $-\text{CH}_3$ or $-\text{O}-\text{CH}_3$.

6. The compound of claim 4 wherein at least one of R_{3a} , R_{3b} or R_{3c} is $-\text{Cl}$ or $-\text{CF}_3$.

7. The compound of claim 1 wherein $-\text{L}_3-\text{Q}$ and $-\text{L}_1-\text{J}$ are the same, where Q and J are an aromatic carbocyclic ring selected from the group consisting of phenyl, substituted phenyl, naphthyl and substituted naphthyl.

8. The compound of claim 1 wherein W comprises an amine, amide, alcohol, carboxylic acid, ether, ester, guanidine or urea, more than one of the foregoing, or a combination of the foregoing.

9. The compound of claim 1 wherein W is



R_4 is

NH ,

O ,

CH_2 , provided that R_5 comprises N or O,

C_6H_5 , provided that R_5 comprises N or O,

$\text{N}(\text{CH}_2)_z$ where $\text{N}(\text{CH}_2)_z$ together with R_5 forms a ring,

$\text{N}((\text{CH}_2)_y-\text{CH}_3)$,

$\text{NH}-\text{C}(=\text{O})$,

$\text{NH}-\text{C}(=\text{O})-\text{NH}$,

$\text{C}(=\text{O})$,

$\text{C}(=\text{O})-\text{NH}$,

$C(=O)-O$, or

$O-C(=O)$;

R_5 is

NH_2 ,

OH ,

CH_3 , provided that R_4 comprises N or O,

$NH-(CH_2)_z$ where $NH-(CH_2)_z$ together with R_4 forms a ring,

$NH-(CH_2)_y-CH_3$,

$N(-(CH_2)_y-CH_3)_2$,

$NH-(CH_2)_z-NH_2$,

$NH-(CH_2)_z-NH-(CH_2)_y-CH_3$,

$NH-(CH_2)_z-N-((CH_2)_y-CH_3)_2$,

$N(-(CH_2)_y-CH_3)-C(=NH)-NH_2$,

$N(-(CH_2)_y-CH_3)-C(=N((CH_2)_y-CH_3))-NH_2$,

$NH-C(=NH)-NH_2$,

$NH-C(=N((CH_2)_y-CH_3))-NH_2$,

$N(-(CH_2)_y-CH_3)-(CH_2)_z-NH(CH_2)_y-CH_3$,

$N(-(CH_2)_y-CH_3)-(CH_2)_z-N((CH_2)_y-CH_3)_2$,

$N(-(CH_2)_y-CH_3)-C(=N((CH_2)_y-CH_3))-NH(CH_2)_y-CH_3$,

$NH-C(=N((CH_2)_y-CH_3))-NH-(CH_2)_y-CH_3$,

$N(-(CH_2)_y-CH_3)-C(=NH)-NH(CH_2)_y-CH_3$,

$NH-C(=N((CH_2)_y-CH_3))-N((CH_2)_y-CH_3)_2$,

$N(-(CH_2)_y-CH_3)-C(=NH)-N((CH_2)_y-CH_3)_2$,

$NH-C(=O)-(CH_2)_y-NH_2$,


$O-(CH_2)_y-CH_3$,

SO_2-NH_2 ,


$SO_2-NH-(CH_2)_y-CH_3$,

$SO_2-N(-(CH_2)_y-CH_3)_2$,


$SO_2-(CH_2)_y-CH_3$,




where one or more of positions 1 to 5 are a heteroatom selected from N for position 1 and S, O or NH for positions 2 to 5,




where none, one or two of positions 1 to 5 are a heteroatom selected from N for position 1 and for the position to which R₆ is bound if such position does not comprise C, and otherwise S, O or NH,




where at least one bond between adjacent ring atoms is a double bond, and one or more of positions 1 to 5 are a heteroatom selected from N for position 1 and any double bond position and otherwise S, O or NH for positions 2 to 5, provided that not more than one position is S or O,




where at least one bond between adjacent ring atoms is a double bond, and one or more of positions 1 to 5 are optionally a heteroatom selected from N for position 1, the position to which R₆ is bound if such position does not comprise C, and any double bond position and otherwise S, O or NH for positions 2 to 5, provided that not more than one position is S or O,




where at least one bond between adjacent ring atoms is a double bond, the oxo is bound to a ring carbon, and one or more of the remaining of positions 1 to 5 are optionally a heteroatom selected from N for position 1 and any double bond position and otherwise S, O or NH for positions 2 to 5, provided that not more than one position is S or O,




where at least one bond between adjacent ring atoms is a double bond, the oxo is bound to a ring carbon, and one or more of positions 1 to 5 are optionally a heteroatom selected from N for position 1, the position to which R_6 is bound if such position does not comprise C, and any double bond position and otherwise S, O or NH for positions 2 to 5, provided that not more than one position is S or O,




where one or more of positions 1 to 6 are a heteroatom selected from N for position 1 and S, O or NH for positions 2 to 6,



where none, one or two of positions 1 to 6 are a heteroatom selected from N for position 1 and the position to which R_6 is bound if such position does not comprise C, and otherwise S, O or NH,




where at least one bond between adjacent ring atoms is a double bond, and one or more of positions 1 to 6 are a heteroatom selected from N for position 1 and any double bond position and otherwise S, O or NH for positions 2 to 6, provided that not more than two positions are S or O,




where at least one bond between adjacent ring atoms is a double bond, and one or more of positions 1 to 6 are optionally a heteroatom selected from N for position 1, the position to which R_6 is bound if such position does not comprise C, and any double bond position and

otherwise S, O or NH for positions 2 to 6, provided that not more than two positions are S or O,



where at least one bond between adjacent ring atoms

is a double bond, the oxo is bound to a ring carbon, and one or more of the remaining of positions 1 to 6 are optionally a heteroatom selected from N for position 1 and any double bond position and otherwise S, O or NH for positions 2 to 6, provided that not more than two positions are S or O, or



where at least one bond between adjacent ring atoms

is a double bond, the oxo is bound to a ring carbon, and one or more of positions 1 to 6 are optionally a heteroatom selected from N for position 1, the position to which R_6 is bound if such position does not comprise C, and any double bond position and otherwise S, O or NH for positions 2 to 6, provided that not more than two positions are S or O;

R_8 is hydroxyl, $(CH_2)_y-CH_3$, $(CH_2)_y-NH_2$, $NH-(CH_2)_y-CH_3$ or $N(-(CH_2)_y-CH_3)_2$;

t is an index value from 0 to 5;


z is an index value from 1 to 6; and

5 y is in each instance independently an index value from 0 to 5;

provided that, any NH or NH_2 in the foregoing may be substituted by N-Prg or NH-Prg, respectively, where each Prg is independently an amine protecting group.

10. The compound of claim 9 wherein each Prg is independently acetyl, adamantyloxy, benzoyl, benzyl, benzyloxycarbonyl, t-butoxycarbonyl, mesitylene-2-sulfonyl, 4-methoxy-2,3,6-trimethyl-benzenesulfonyl, 2,2,4,6,7-pentamethyldihydrobenzofurane-5-sulfonyl, 2,2,5,7,8-pentamethylchromane-6-sulfonyl, 9-fluorenylmethyloxycarbonyl, or tosyl.

11. The compound of claim 1 of a formula:



wherein:

5 R_7 is H or =O;

R_8 is hydrogen or $N(R_{9a}R_{9b})$;

R_{9a} and R_{9b} are each independently each independently hydrogen, acetyl, methyl, ethyl, propyl, isopropyl, butyl, pentyl, hexyl, isobutyl, benzyl, benzoyl, hexanoyl, propionyl, butanoyl, pentanoyl, heptanoyl, cyclopropyl, cyclopropylmethyl, cyclobutyl, cyclobutylmethyl, cyclohexyl, cyclohexylmethyl, or polyethylene glycol;


v is in each instance independently an index value from 0 to 2; and

y is in each instance independently an index value from 0 to 5;

wherein if R_8 is not hydrogen, the adjacent carbon atom marked with an asterisk can have any stereochemical configuration.

12. The compound of claim 11 wherein the polyethylene glycol has a formula molecular weight of between 100 and 50,000.

5 13. The compound of claim 1 wherein one of R_{2a} and R_{2b} is



and the remaining of R_{2a} and R_{2b} and both R_{1a} and R_{1b} are hydrogen.

14. The compound of claim 1 wherein W comprises a single amine.

10

15. The compound of claim 1 wherein W comprises a single N and a single O.

16. The compound of claim 1 wherein W comprises and $-NH-C(=NH)-NH_2$.

15

17. The compound of claim 1, wherein the compound is

N-{3-[(2S,5R)-1-((R)-2-amino-3-naphthalen-2-yl-propionyl)-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;

20

N-{3-[(2S,5R)-5-benzyl-1-((R)-2-dimethylamino-3-naphthalen-2-yl-propionyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;

N-{3-[(2S,5R)-5-benzyl-1-(4'-chloro-biphenyl-2-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;

25

N-{3-[(2S,5R)-5-benzyl-1-(4'-chloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;

30

N-{3-[(2S,5R)-5-benzyl-1-(4'-chloro-biphenyl-4-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;

N-{3-[(2S,5R)-5-benzyl-1-[2-(4'-chloro-biphenyl-2-yl)-acetyl]-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;

35

N-{3-[(2S,5R)-5-benzyl-1-(2',4'-dichloro-biphenyl-2-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;

N-{3-[(2S,5R)-5-benzyl-1-(3',4'-dichloro-biphenyl-2-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;

40

- N-{3-[(2S,5R)-5-benzyl-1-(3',5'-dichloro-biphenyl-2-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 5 N-{3-[(2S,5R)-1-((R)-2-amino-3-naphthalen-2-yl-propyl)-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- N-{3-[(2S,5R)-5-benzyl-1-(2',4'-dichloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 10 N-{3-[(2S,5R)-5-benzyl-1-(2',4'-dichloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- N-{3-[(2S,5R)-5-benzyl-1-(3',4'-dichloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 15 N-{3-[(2S,5R)-5-benzyl-1-(3',5'-dichloro-biphenyl-3-carbonyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- N-{3-[(2S,5R)-5-benzyl-1-(3-iodo-benzoyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 20 N-{3-[(2S,5R)-5-benzyl-1-(2',4'-dichloro-biphenyl-3-ylmethyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- N-{3-[(2S,5R)-5-benzyl-1-(4'-chloro-biphenyl-2-ylmethyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 25 N-{3-[(2S,5R)-5-benzyl-1-(4'-chloro-biphenyl-3-ylmethyl)-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 30 N-{3-[(2S,5R)-5-benzyl-1-biphenyl-2-ylmethyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- N-{3-[(2S,5R)-1-[(R)-2-amino-3-(2,4-dimethyl-phenyl)-propyl]-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 35 N-{3-[(2S,5R)-1-[(R)-2-amino-3-naphthalen-2-yl-propionyl]-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 40 N-{3-[(2R,5R)-1-[(R)-2-amino-3-naphthalen-2-yl-propionyl]-5-benzyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- N-{3-[(2S,5R)-1-[(R)-2-amino-3-(2,4-dimethyl-phenyl)-propyl]-5-cyclohexylmethyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 45 N-{3-[(2S,5R)-1-[(R)-2-amino-3-(2,4-dichloro-phenyl)-propyl]-5-cyclohexylmethyl-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- N-{3-[(2S,5R)-5-cyclohexylmethyl-1-[(R)-3-(2,4-dichloro-phenyl)-2-dimethylamino-propyl]-4-(2-naphthalen-2-yl-acetyl)-piperazin-2-yl]-propyl}-guanidine;
- 50 (R)-2-amino-1-[(2R,5S)-4-[(R)-2-amino-3-(2,4-dimethyl-phenyl)-propionyl]-5-[3-(2-amino-ethylamino)-propyl]-2-cyclohexylmethyl-piperazin-1-yl]-3-(2,4-dimethyl-phenyl)-propan-1-one;
- 55 (R)-2-amino-1-[(2R,5S)-4-[(R)-2-amino-3-(2,4-dichloro-phenyl)-propionyl]-5-[3-(2-amino-ethylamino)-propyl]-2-cyclohexylmethyl-piperazin-1-yl]-3-(2,4-dichloro-phenyl)-propan-1-one;
- (R)-2-amino-1-[(2R,5S)-5-[3-(2-amino-ethylamino)-propyl]-4-[(R)-2-amino-3-naphthalen-2-yl-


propionyl)-2-cyclohexylmethyl-piperazin-1-yl]-3-naphthalen-2-yl-propan-1-one;

(R)-2-amino-1-[(2S,5R)-2-[4-(2-amino-ethylamino)-butyl]-4-((R)-2-amino-3-naphthalen-2-yl-propionyl)-5-cyclohexylmethyl-piperazin-1-yl]-3-naphthalen-2-yl-propan-1-one; or

(R)-2-amino-1-[(2S,5R)-4-[(R)-2-amino-3-(2,4-dichloro-phenyl)-propionyl]-2-[4-(2-amino-ethylamino)-butyl]-5-cyclohexylmethyl-piperazin-1-yl]-3-(2,4-dichloro-phenyl)-propan-1-one,

or a pharmaceutically acceptable salt thereof.

18. A compound having the formula of structure II:



or an enantiomer, stereoisomer or diastereoisomer thereof, or a pharmaceutically acceptable salt thereof,

wherein

each J is independently a ring structure selected from the group consisting of substituted or unsubstituted aromatic carbocyclic rings, substituted or unsubstituted non-aromatic carbocyclic rings, substituted or unsubstituted aromatic fused carbobicyclic ring groups, two substituted or unsubstituted aromatic carbocyclic rings wherein the rings are joined by a bond, -CH₂-, or -O-, and substituted or unsubstituted aromatic fused heterobicyclic ring groups, wherein in each instance the rings include 5 or 6 ring atoms;


W is a heteroatom unit with at least one cationic center, hydrogen bond donor or hydrogen bond acceptor wherein at least one heteroatom is nitrogen or oxygen;

L₁ is a bond or a linker unit comprising from one to eight backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

L₂ is a bond or -(CH₂)_z-;


L₃ is a bond or a linker unit comprising from one to eight backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

R_{1a}, R_{1b}, R_{2a} and R_{2b} are selected such that:



the remaining of R_{2a} and R_{2b} is hydrogen and R_{1a} and R_{1b} are each hydrogen, R_{1a} and R_{1b} together form $=O$, or one of R_{1a} and R_{1b} is a C_1 to C_6 aliphatic linear or branched chain and the remaining of R_{1a} and R_{2b} is hydrogen,

5 or R_{2a} and R_{2b} are each hydrogen



and the remaining of R_{1a} and R_{1b} is hydrogen;





X is CH_2 , $C=O$ or $C=S$;








z is an index value from 1 to 6; and

10 y is an index value from 0 to 5;

wherein the carbon atom marked with an asterisk can have any stereochemical configuration.

19. The compound of claim 18 wherein each J is independently






unsubstituted or substituted with one or more ring substituents.

20. The compound of claim 19 wherein J is substituted with one or more ring
5 substituents independently selected from the group consisting of hydroxyl, halogen,
sulfonamide, alkyl, -O-alkyl, aryl, and -O-aryl.

21. The compound of claim 18 wherein -L₃-J and -L₁-J are the same.

- 10 22. A pharmaceutical composition comprising a compound of claim 1 and a
pharmaceutically acceptable carrier.

23. A use of a compound of structure I



or an enantiomer, stereoisomer or diastereoisomer thereof, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for the treatment of a condition responsive to changes in melanocortin receptor function in a human or non-human mammal, wherein

J is a ring structure selected from the group consisting of substituted or unsubstituted aromatic carbocyclic rings, substituted or unsubstituted non-aromatic carbocyclic rings, substituted or unsubstituted aromatic fused carbobicyclic ring groups, two substituted or unsubstituted aromatic carbocyclic rings wherein the rings are joined by a bond, $-\text{CH}_2-$, or $-\text{O}-$, and substituted or unsubstituted aromatic fused heterobicyclic ring groups, wherein in each instance the rings include 5 or 6 ring atoms;

W is a heteroatom unit with at least one cationic center, hydrogen bond donor or hydrogen bond acceptor wherein at least one heteroatom is nitrogen or oxygen;


Q is an aromatic carbocyclic ring selected from the group consisting of phenyl, substituted phenyl, naphthyl and substituted naphthyl;

L_1 is a bond or a linker unit comprising from one to eight backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

L_2 is a bond or $-(\text{CH}_2)_z-$;

L_3 is a bond or a linker unit comprising from one to nine backbone atoms selected from the group consisting of carbon, sulfur, oxygen or nitrogen;

R_{1a} , R_{1b} , R_{2a} and R_{2b} are selected such that:




the remaining of R_{2a} and R_{2b} is hydrogen and R_{1a} and R_{1b} are each hydrogen, R_{1a} and R_{1b}

together form =O, or one of R_{1a} and R_{1b} is a C₁ to C₆ aliphatic linear or branched chain and the remaining of R_{1a} and R_{2b} is hydrogen,

or R_{2a} and R_{2b} are each hydrogen

and one of R_{1a} and R_{1b} are



5 and the remaining of R_{1a} and R_{1b} is hydrogen;

X is CH₂, C=O or C=S;

z is an index value from 1 to 6; and

y is an index value from 0 to 5;

wherein the carbon atom marked with an asterisk can have any stereochemical

10 configuration.

24. The use of claim 23, wherein the condition responsive to changes in melanocortin receptor function is selected from the group consisting of male sexual dysfunction, female sexual dysfunction, an eating disorder, above-optimal body weight, obesity, below-optimal body weight and cachexia.

15